Out of the pot and into the fire: Explaining the vulnerability of an endangered small headwater stream fish to black-bass *Micropterus* spp. invasion

Ellender, B. R., Weyl, O. L. F., Alexander, M. E., Luger, A. M., Nagelkerke, L. A. J., & Woodford, D. J.

This article is made publically available in the institutional repository of Wageningen University and Research, under article 25fa of the Dutch Copyright Act, also known as the Amendment Taverne.

Article 25fa states that the author of a short scientific work funded either wholly or partially by Dutch public funds is entitled to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

For questions regarding the public availability of this article, please contact openscience.library@wur.nl.

Please cite this publication as follows:

Ellender, B. R., Weyl, O. L. F., Alexander, M. E., Luger, A. M., Nagelkerke, L. A. J., & Woodford, D. J. (2018). Out of the pot and into the fire: Explaining the vulnerability of an endangered small headwater stream fish to black-bass *Micropterus* spp. invasion. *Journal of Fish Biology*, 92(4), 1035-1050.
<https://doi.org/10.1111/jfb.13562>

Out of the pot and into the fire: Explaining the vulnerability of an endangered small headwater stream fish to black-bass *Micropterus* spp. invasion

B. R. ELLENDER*†‡, O. L. F. WEYL*†§, M. E. ALEXANDER†||, A. M. LUGER†¶, L. A. J. NAGELKERKE¶ AND D. J. WOODFORD†**

*Department of Ichthyology and Fisheries Science, P.O. Box 96, Grahamstown, 6140, South Africa, †Centre for Invasion Biology, South African Institute for Aquatic Biodiversity, Private Bag 1015, Grahamstown, 6140, South Africa, §DST/NRF Research Chair in Inland Fisheries and Freshwater Ecology, South African Institute for Aquatic Biodiversity, Grahamstown, South Africa, ||Institute of Biomedical and Environmental Health Research (IBEHR), School of Science and Sport, University of West Scotland, High Street, Paisley, PA1 2BE, ¶Aquaculture and Fisheries Group, Wageningen University, Wageningen, The Netherlands and **Centre for Invasion Biology, School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Private Bag 3, Wits, 2050, South Africa

(Received 26 August 2017, Accepted 14 January 2018)

Introduced predatory fishes have had consistently severe consequences for native fishes in stream environments around the world, although the drivers of these effects are often unclear. In the Swartkops River headwaters in South Africa, native Eastern Cape redfin *Pseudobarbus afer* were always absent from sites occupied by non-native black basses *Micropterus salmoides* and *Micropterus dolomieu*, but generally co-occurred with the native predators *Anguilla marmorata* and *Anguilla mossambica*. A natural experiment provided by flood-mediated recolonization of black-bass occupied sites by *P. afer* demonstrated depletion in black-bass invaded sites. Field behavioural observations of *P. afer* indicated that they foraged among benthic cover during the day, but suspended in open water at night. As the nocturnal *A. marmorata* and *A. mossambica* foraged actively within structural cover at night and *M. dolomieu* and *M. salmoides* are diurnal or crepuscular predators, *P. afer* is thus optimized to avoid predation by native anguillid predators and not the functionally unique predatory black basses. The integration of distributional, temporal population dynamics and behavioural data suggests that the severe effects of *Micropterus* spp. are probably a consequence of prey naïveté and behaviour evolved to evade native predators.

© 2018 The Fisheries Society of the British Isles

Key words: behaviour; endangered fish; evolutionary response; *Micropterus*; naïveté; *Pseudobarbus afer*.

INTRODUCTION

Fish communities in stream headwaters are among the most adversely affected by non-native fish invasions because they often comprise unique assemblages of

‡Author to whom correspondence should be addressed. Tel.: +27 (0)46 603 5823; email: bru.ellender@gmail.com

small-bodied specialist fishes (Meyer *et al.*, 2007). This is particularly apparent in Mediterranean-climate region streams, where the introduced species diversity is often higher than that of natives (Marr *et al.*, 2010), and native headwater fish are typically range restricted and isolated in headwater refugia (Henriques *et al.*, 2010; Marr *et al.*, 2010; Weyl *et al.*, 2013).

The effects of introduced black-bass *Micropterus* Lacépède 1802 species on small native fishes appear to be particularly severe. In the Raia tributary of the Tagus River in Portugal, for example, the presence of largemouth bass *Micropterus salmoides* (Lacépède 1802) was the best biotic predictor of native fish species assemblage structure, with small cyprinids being absent from reaches invaded by *M. salmoides* (Godinho & Ferreira, 2000). In the intermittently flowing streams of the Guadiana River system, predation by non-native fish was a major contributing factor to the decline of the small cyprinid *Anaecypris hispanica* (Steindacher 1866), which is consequently at high risk of extinction (Collares-Pereira *et al.*, 1999). In an assessment of 41 tributaries of the Olifants–Doorn River system in South Africa, van der Walt *et al.*, (2016) demonstrated that *Micropterus dolomieu* Lacépède 1802 and *Micropterus punctulatus* (Rafinesque 1819) had consistently extirpated small-bodied cyprinid minnows from invaded reaches. Similarly, since the introduction of non-native species to the Cosumnes River basin, California (U.S.A.), there has been a gradual disappearance of native fish. In some areas invasive redeye bass *Micropterus coosae* (Hubbs & Bailey 1940) represented >90% of fish collected (Moyle & Williams, 1990; Moyle *et al.*, 2003).

In Mediterranean-climate region streams in South Africa the endpoint of invasions by predatory fishes into headwater streams has been relatively well documented and is consistent with invasions of similar streams in other regions globally (Ellender *et al.*, 2017). Invasions of streams in South Africa typically lead to complete extirpation of native fishes in invaded zones (Ellender *et al.*, 2011; Weyl *et al.*, 2013; van der Walt *et al.*, 2016). Consequently, headwater stream fishes are increasingly being isolated in small fragmented headwater refuges due to invasion by non-native fishes (Woodford *et al.*, 2005; Clark *et al.*, 2009; van der Walt *et al.*, 2016). One such case is the Blindekloof River, a Swartkops River headwater tributary, where the Eastern Cape redbfin *Pseudobarbus afer* (Peters 1864) was isolated in headwater refugia and extirpated from invaded reaches further downstream by black-bass, *M. salmoides* and *M. dolomieu*, predation (Ellender *et al.*, 2011). Although the consequences of predatory fish invasions of headwater streams have been relatively well documented, the drivers responsible for the magnitude of the effects on headwater stream fishes remain unclear. This is further complicated by the context-dependent nature of perturbations, with very few rules on specific effects resulting from non-native species introductions (Ricciardi & Atkinson, 2004). While there is evidence to suggest that distinctiveness (Ricciardi & Atkinson, 2004), prey naïveté (a lack of defence toward a functionally unique predator) (Cox & Lima, 2006) and behaviour (Kadye & Booth 2014) are contributory factors, the ecological implications of biological invasions remains the most poorly understood aspect of the invasion process (Parker *et al.*, 1999; Kulhanek *et al.*, 2011).

This study integrates abundance, distributions and behavioural data to document invasion at a system scale, quantify effects by documenting depletion of native fishes in invaded v. non-invaded stream reaches and uses behavioural observations to infer naïveté as a mechanism for the observed low tolerance of *P. afer*, a small, endangered, headwater stream fish, to non-native predatory fish invasions.

FIG. 1. The position of the Swartkops River system headwater streams, sites sampled (●) within the Groendal Wilderness Area, Eastern Cape, South Africa. The position of the gauging weir (M1H004) on the Elands River and the weather station (M1E001) at Groendal Dam are also shown.

MATERIALS & METHODS

STUDY SITE

The study was conducted in the headwaters of the Swartkops River system, a relatively short coastal river system that flows for 110 km from its source in the Grootwinterhoek Mountains (1000 m a.s.l.) before the river enters the Indian Ocean near the city of Port Elizabeth (Fig. 1). This study focussed on six first- and second-order (Strahler, 1957) headwater tributary streams (Blindekloof, Chaseskloof, Fernkloof, Nounekloof, Vyeboomkloof and Waterkloof) of the Kwa-Zunga (third-order stream) tributary of the Swartkops River, South Africa. The headwater

streams are free flowing, but the main stem Kwa-Zunga River was dammed in 1933 to create the 96 ha Groendal Dam (Fig. 1). The stream catchments are otherwise relatively pristine and unaffected by human influence because they are located in a protected area, the Groendal Wilderness Area, which forms part of the Baviaanskloof Mega-Reserve cluster.

SURVEY (1): DISTRIBUTION

To investigate the factors influencing distribution of native and non-native fishes at a system scale, 69 sites were sampled by backpack electrofishing and snorkel surveys during 2010. In order to obtain as comprehensive cover as possible, sampling sites were selected to cover the six headwater streams as well as the mainstream environment. At each sampling site, temperature, conductivity and pH were measured using a Hanna HI98129 Combo pH and electrical conductivity meter (HANNA Instruments Inc.; www.hannainst.com). Turbidity was measured using a Hanna HI 98703 turbidimeter (HANNA Instruments Inc.). To estimate pool volume and habitat diversity within each pool, the length of the pool was measured, followed by four to six equally spaced width measurements. On each width transect, three depths were measured, the outer two were each 0.2 m from the left and right-hand stream bank and the third measurement taken midstream. At each depth measurement, the habitat type was recorded. Canopy cover was estimated as a percentage of total cover and bankside vegetation type was recorded.

At each site the choice of sampling method (snorkel survey or backpack electrofishing) depended on the dimensions of the sampling site. Snorkel surveys were conducted in streams where the depth of the sampled pools was >1 m, using a two-pass zigzag method as described in Ellender *et al.* (2011). Pass one was initiated from the tail end of the pool, with the observer swimming upstream, zigzagging to cover as much of the pool as possible. Pass two was a repeat of pass one, but in a downstream direction. During each pass, all fish seen were identified to species level and counted. Sites where the average depth of the stream was <1 m were sampled by single pass electrofishing. Electrofishing was conducted using a Samus 725G backpack electrofisher (SAMUS; www.electrofisher.net), attached to a 12 V battery with settings standardized with a duration of 0.3 ms and a frequency of 80 Hz. The pass was conducted from the downstream side (tail) of the pool in an upstream direction, covering the entire length of each pool. Upon completion of the pass, fish were identified to species level, measured, counted and released alive. Block nets were not used as pools were mostly isolated from each other. To ensure consistency and standardized methods of data collection, all snorkelling and electrofishing were undertaken by the same individual.

The necessity for using a multi-method approach for overall fish distribution surveys resulted in a lack of standardized catch data between streams. Therefore, for overall distribution data only presence-absence data were analysed using CANOCO 4.5 (ter Braak & Šmilauer, 2002). For analyses, data were separated into two matrices. The first consisted of the site specific presence-absence data and the second were the environmental data. Environmental data were standardized with z transformation by transforming the original distribution to a mean of 0 and S.D. = 1. Preliminary analyses using detrended correspondence analysis (DCA), indicated gradient lengths of >2 on the first axis, therefore unimodal ordination methods were chosen for analysis of overall distribution data in the form of correspondence analysis (CA, species data only) and canonical correspondence analysis (CCA, species-environmental relation) (ter Braak, 1995). Initially, eight environmental variables were used in the CCA: canopy cover, altitude, instream barriers (natural waterfalls), maximum depth, habitat diversity, surface area, temperature and turbidity. Environmental variables with variance inflation factors (VIF) > 10 were then removed from further analyses due to multicollinearity with other variables (ter Braak & Šmilauer, 1998). A forward step-wise procedure was then used to determine the best predictor variables. The significance of these variables' contribution to the ordination was tested using 999 Monte-Carlo simulations at the $P < 0.05$ level. To test the statistical significance of the selected environmental variables on the species presence-absence model, Monte-Carlo permutation tests were used.

SURVEY (2): MONTHLY MONITORING

Baseline distributions data indicated that at a system scale, no native fishes co-occurred with black basses. A major flood followed the baseline distribution surveys (Ellender & Weyl,

2015). Post-flood exploratory surveys indicated that on the Blindekloof River, a representative Swartkops River system headwater tributary, native fishes were distributed throughout the stream and now inhabited sites that were invaded by black basses. This provided an ideal opportunity to use this incursion of native fishes into black-bass invaded reaches as a natural experiment to compare their abundance, distribution and population structure in the black-bass invaded stream reaches to those in the non-invaded (control) stream reaches. To do this, snorkel surveys were conducted monthly (November 2011–April 2012) to assess for the distribution range, relative abundance and population structure of native fishes.

For subsequent analysis, the Blindekloof River was divided into 15 sites that represented three invasion states according to the presence–absence of black basses: sites 1–4 represented the mixed zone (no black basses, but other non-native species present); sites 5–9 the invaded zone (black basses present); sites 10–15 the non-invaded zone (only native species present). Despite the episodic nature of the Blindekloof River, pools are fed by groundwater and so surface area and volume remained fairly constant throughout the study. It was therefore assumed that effort at each monitoring site was also constant and comparisons between snorkel counts would be sufficient to reflect any temporal abundance changes.

Snorkel surveys were conducted using the zigzag method described for survey (1), except that, in addition to species-level identification and enumeration during each pass, fish were also categorized into one of four predetermined total length (L_T) classes: 0–15, 15–30, 30–60 and >60 mm. To estimate L_T , the snorkeler carried a slate with a ruler attached to it, where the snout and tail of the fish are aligned with the ruler (Cunjak & Power, 1986). To avoid measuring the same individual twice, length was estimated only on the first snorkel pass.

Two native predatory eels are found in the headwaters of the Swartkops River, the giant mottled eel *Anguilla marmorata* Quoy & Gaimard 1824 and the longfin eel *Anguilla mossambica* (Peters 1852). Both eel species are nocturnal and in order to get a representative sample of the relative abundance of native predators it was necessary to undertake night snorkelling. Identical methods were used during the day and at night, except that fish lengths were not estimated at night. A 400 lm SL4 eLED dive torch (Underwater Kinetics; www.uwkinetics.com) was used for night observations. For those sites sampled during the day and at night, snorkelling was undertaken during the same 24 h period (Thurrow & Schill, 1996). Night snorkelling commenced a minimum of 45 min after sunset in all months except April 2012 due to equipment failure).

Data from the 6 month monitoring of the 15 monitoring sites were used to assess for temporal differences in abundance of *P. afer* irrespective of size class (analysis 1), young of year (YOY; ≤ 30 mm L_T ; analysis 2) and adults (≥ 30 mm L_T ; analysis 3). For size-class comparisons, only first-pass snorkelling data were used. First-pass data were considered representative as there was a highly significant correlation between *P. afer* abundance from the first and second passes (Spearman $r = 0.908$, $P < 0.001$). These data were analysed using linear mixed models in SPSS 16.0 (IBM; www.ibm.com). Abundance analyses were undertaken using restricted maximum likelihood models in SPSS. Fixed effects for all three analyses were specified as stream zone (mixed zone, invaded zone and non-invaded zone), sample months (1–6) were added as a covariate and site (1–15) was included as a random effect. Analysis 1 used overall *P. afer* abundance data (6 months). Analysis 2 used the number of *P. afer* young of the year (YOY). Analysis 3 used adult *P. afer* abundance. Main treatment effects were analysed using least significant difference (LSD) pair-wise comparisons.

To test whether native eels were evenly distributed throughout the study reach (and therefore apply similar predation pressure throughout the study reach), the frequency of occurrence (% of sites within each stream zone where *A. mossambica* and *A. marmorata* were recorded) was tested for uniform distribution between stream zones using a χ^2 -test of independence.

SURVEY (3): BEHAVIOUR

Snorkel survey observations (BRE) of the diurnal and nocturnal behaviour of native fishes indicated that this might provide some insight into their vulnerability to introduced predators. To investigate for diurnal differences in fish behaviour a survey of 32 sites on the Blindekloof River was undertaken during the day and at night in November 2011 using identical methods to survey 2. To support behavioural observations using abundance data, day and night counts of *P. afer* were compared using a paired *t*-test (RStudio Team, 2015).

FIG. 2. Canonical correspondence ordination biplot of species and environmental variables for the overall distribution data of fishes from headwater tributaries of the Swartkops River within the Groendal Wilderness Area. Species: *Pseudobarbus afer*, *Enteromius pallidus*, *Sandelia capensis*, *Tilapia sparrmanii*, *Glossogobius callidus*, *Clarias gariepinus*, *Micropterus dolomieu*, and *Micropterus salmoides*.

All field research was reviewed by and conducted according to the ethical code for the field sampling of fishes set out by the South African Institute for Aquatic Biodiversity.

RESULTS

DISTRIBUTION

Eight fish species were recorded from the surveyed streams, four natives, *P. afer*, kurper *Sandelia capensis* (Cuvier 1829), goldie barb *Enteromius pallidus* (Smith 1841), river goby *Glossogobius callidus* (Smith 1937), and four non-natives *M. salmoides*, *M. dolomieu*, banded tilapia *Tilapia sparrmanii* Smith, 1840 and African sharp-tooth catfish *Clarias gariepinus* (Burchell 1822). Of the 29 sites where *P. afer* occurred, they co-occurred with all species other than black basses (>60% with *S. capensis*; <11% with *G. callidus*, *E. pallidus*, *C. gariepinus* and *T. sparrmanii*). In the non-invaded stream reaches, *P. afer* were widespread, inhabiting uninhabited segments of five of the six surveyed streams (Blindekloof, Chaseskloof, Fernkloof, Nounekkloof and Waterkloof).

The CCA identified three distinct assemblages (native: *P. afer* and *S. capensis*; mixed: *E. pallidus*, *G. callidus*, *C. gariepinus* and *T. sparrmanii*; black basses: *M. dolomieu* and *M. salmoides*) explaining 98% of the overall species-environment relation on the first two axes (Fig. 2). The CCA's explanatory variables (altitude, surface area and

TABLE I. Summary statistics for canonical correspondence analysis (CCA, species and environmental data) showing the contribution of fish species and environmental variation to overall fish species assemblages for the distribution data from the headwater tributary streams of the Swartkops River within the Groendal Wilderness Area

Statistic	Axis 1	Axis 2
Canonical correspondence analysis		
Eigenvalue	0.50	0.22
Species–environment correlation	0.79	0.51
Cumulative percentage variance of species data	18.9	27.0
Cumulative percentage variance of species-environment relation	68.9	98.0
Weighted correlations		
Altitude	−0.31	0.65
Surface Area	0.07	0.94
Temperature	0.19	1.0
Total inertia		2.68
Sum of all canonical eigenvalues (full CCA)		0.73

Only environmental variables with a significant contribution to assemblage variation are listed.

temperature) significantly accounted for 27.4% of the species-environmental variance (Monte Carlo permutation $P < 0.05$, of both the first axis and trace) (Table I). The native assemblage had high association with the mixed assemblage, but no association with the black-bass assemblage.

CC axis 1 suggested an association of black basses with altitude and surface area whereas *P. afer* and *S. capensis* were consistent throughout a range of altitudes and smaller surface-area environments (Fig. 2). The CC axis 2 suggested a strong positive relationship between species in the mixed assemblage with pool surface area and higher temperatures and *P. afer* with lower temperatures (Fig. 2).

MONTHLY MONITORING

The overall abundance, relative abundance and distribution of fishes from the Blindekloof River monitoring sites is summarized in Supporting Information Table SI. Nine species were recorded, six native (*P. afer*, *S. capensis*, *E. pallidus*, *G. callidus*, *A. marmorata*, *A. mossambica*) and three non-natives (*M. dolomieu*, *T. sparrmanii*, *C. gariepinus*). The mixed zone was characterized by an assemblage consisting of the native species *P. afer*, *E. pallidus*, *G. callidus*, *S. capensis*, *A. mossambica* and *A. marmorata* and the non-native species *T. sparrmanii* and *C. gariepinus*. The invaded zone consisted of five sampling sites, three of which were inhabited by *M. dolomieu* and two sites in between by native fishes. In the non-invaded zone only *P. afer*, *S. capensis*, *G. callidus*, *A. mossambica* and *A. marmorata* were recorded.

The frequency of occurrence of the native predators *A. mossambica* ($n = 11$ sites) and *A. marmorata* ($n = 8$ sites) was not significantly dependent on stream zone (*A. marmorata*: χ^2 -test of independence: $\chi^2 = 2.81$, d.f. = 10, $P > 0.05$; *A. mossambica*: $\chi^2 = 0.61$, d.f. = 10, $P > 0.05$). Overall *P. afer* abundances did not differ significantly between stream zones ($F_{2,23.96} = 3.13$, $P > 0.05$) but followed a downstream gradient and were highest in the non-invaded zone (mean $\pm$ S.E., 108.6 ± 36.8

FIG. 3. Mean $\pm$ S.E. abundance (N) of the Eastern Cape redfin *Pseudobarbus afer* for: (a) all age groups (analysis 1); (b) young-of-the-year (analysis 2); (c) adult fish (analysis 3). White, grey and black bars represent non-invaded, mixed and invaded zones, respectively.

fish 100 m^{-2}), lower in the mixed zone (25.1 ± 8.5 fish 100 m^{-2}) and lowest in the invaded zone (19.7 ± 8.5 fish 100 m^{-2}) [Fig. 3(a)]. Overall *P. afer* abundances differed significantly by sample month ($F_{1,34.56} = 33.55$, $P < 0.05$), but no significant interactions between sample month and stream zone were observed. Abundance in all three stream zones increased throughout the monitoring period, with lowest abundance in November 2011 (non-invaded zone: 149.9 ± 50.3 fish 100 m^{-2} ; invaded zone: 20.4 ± 9.0 fish 100 m^{-2} ; mixed zone: 22.9 ± 7.4 fish 100 m^{-2}) and highest abundances in January 2012 (non-invaded zone: 212.1 ± 56.6 fish 100 m^{-2} ; invaded zone: 52.7 ± 10.2 fish 100 m^{-2} ; mixed zone: 61.7 ± 11.7 fish 100 m^{-2}) [Fig. 3(a)].

Changes in YOY abundance were significantly explained by stream zone ($F_{2,16.61} = 5.08$, $P < 0.05$) and an interaction between stream zone and sample month ($F_{2,28.48} = 3.72$, $P < 0.05$). The invaded (21.0 ± 7.2 fish 100 m^{-2} ; $P < 0.05$) and mixed (25.1 ± 6.9 fish 100 m^{-2} ; $P > 0.05$) zones were significantly lower than the non-invaded zone (96.2 ± 22.9 fish 100 m^{-2} ; $P > 0.05$), but not significantly different from each other [Fig. 3(b)]. Young of the year abundances were low during November 2011 (non-invaded zone: 103.5 ± 38.9 fish 100 m^{-2} ; invaded zone: 17.3 ± 7.2 fish 100 m^{-2} ; mixed zone: 12.4 ± 3.6 fish 100 m^{-2}), increasing monthly and peaking in January

FIG. 4. Mean $\pm$ S.E. abundance (N) of the Eastern Cape redfin *Pseudobarbus afer* in the invaded zone over the 6 month monitoring period in the Blindekloof River, a Swartkops headwater tributary within the Groendal Wilderness Area: (a) within the sites where *Micropterus dolomieu* was consistently recorded ($n = 3$) (—●—, rainfall); (b) sites where *M. dolomieu* was not recorded ($n = 2$).

2012 (non-invaded zone: 187.4 ± 49.0 fish 100 m⁻²; invaded zone: 55.4 ± 10.3 fish 100 m⁻²; mixed zone: 56.1 ± 13.1 fish 100 m⁻²) [Fig. 3(b)].

Analysis of adult *P. afer* abundances indicated that stream zone ($F_{2,40.05} = 6.51$, $P < 0.05$) and sample month and stream zone interactions ($F_{2,72} = 5.76$, $P < 0.05$) were significant factors influencing adult abundance [Fig. 3(c)]. Adult *P. afer* abundances differed significantly between zones ($P < 0.05$), with lowest abundances recorded in the invaded (1.6 ± 0.7 fish 100 m⁻²), higher in the mixed (10.0 ± 1.8 fish 100 m⁻²) and highest in the non-invaded zone (39.1 ± 8.0 fish 100 m⁻²). There were no differences in overall *P. afer* abundance by month in the invaded zone ($P < 0.05$). In the non-invaded zone, there was a negative relationship between *P. afer* abundance and sample month, with abundance decreasing from November 2011 (46.4 ± 17.7 fish 100 m⁻²) to April 2012 (14.4 ± 5.3 fish 100 m⁻²), while in the mixed zone, *P. afer* abundance was dynamic, with abundances ranging from the lowest in February 2012 (4.4 ± 2.0 fish 100 m⁻²) to the highest in March 2012 (14.9 ± 4.4 fish 100 m⁻²) [Fig. 3(c)].

At sites within the invaded zone where *M. dolomieu* was consistently recorded, *P. afer* adult abundance decreased over time [Fig. 4(a)] compared with those pools where *M. dolomieu* was not recorded [Fig. 4(b)]. During months of decreasing rainfall (November 2011–January 2012) *P. afer* abundance decreased to non-detectable levels in *M. dolomieu* inhabited sites (November 2011 = 0.9 ± 0.9 fish 100 m⁻²; February = 0 fish). Thereafter, flows following high rainfall in February and March resulted in *P. afer* again dispersing into these sites at very low abundances (0.2 ± 0.1 fish 100 m⁻²).

In sites where *M. dolomieu* was absent, *P. afer* adult abundance was initially high during November 2011 (10.5 ± 1.0 fish 100 m^2), dropped to zero in December 2011 and thereafter fluctuated between January 2012 (2.5 ± 1.7 fish 100 m^2) and April 2012 (4.8 ± 1.5 fish 100 m^2). There was also a lag effect in the dispersal of adults into the invaded zone, with adults appearing 2 months after high rainfall during February and March 2012.

BEHAVIOUR

During the day *P. afer* were observed swimming in the water column and hiding in cracks and under rocks while at night they were predominantly suspended in the water column, well off the bottom of the stream. At night *A. mossambica* and *A. marmorata* were hunting on the bottom of the stream, examining cracks and crevices while they systematically covered a pool and neither species were observed during the day. The majority of surveyed sites (68.8%) had higher *P. afer* night compared with day counts. Day and night counts differed significantly (day: 18.0 ± 4.2 fish 100 m^2 ; night: 27.0 ± 5.6 fish 100 m^2 ; $P < 0.05$).

DISCUSSION

Surveys under base flow conditions demonstrated that reaches invaded by black basses (*M. salmoides* and *M. dolomieu*) were devoid of *P. afer* and other small native fishes, a finding consistent with their presence on other headwater streams (Moyle *et al.*, 2003; Kimberg *et al.*, 2014; van der Walt *et al.*, 2017). By integrating approaches to investigate distribution, temporal population dynamics and behavioural data, the drivers of this observed effect could be further elucidated.

In the headwaters of the Swartkops River system, three distinct assemblages were identified under base-flow conditions (native, mixed and black basses). This is not unusual as biotic (predation, competition) and abiotic features of streams over any spatial scale (stream reach or stream order) typically result in variable species composition of fish assemblages (Angermeier & Winston, 1999; Jackson *et al.*, 2001; Smith & Kraft, 2005). There was considerable overlap between the mixed (*B. pallidus*, *G. callidus*, *T. sparrmanii*, *C. gariepinus*) and native (*P. afer*, *S. capensis*) assemblages, but species from neither assemblage co-occurred with the black-bass assemblage. Associations between the three assemblages and both physical (surface area and altitude) and physico-chemical (temperature) factors were weak, therefore, separation on an environmental gradient did not adequately explain the absence of both the mixed and native assemblage in black-bass invaded zones. The three distinct assemblages are postulated to be controlled by predation, as has been found in other headwater stream environments invaded by black basses (Schlosser, 1987; Godinho & Ferreira, 2000; MacRae & Jackson, 2001).

Piscivory can be a major contributor to the variation and structure of stream-fish assemblages (Jackson *et al.*, 2001; Moyle *et al.*, 2003) as is demonstrated by this study, where the presence of black basses resulted in the absence of all other fishes from invaded zones. Similar results have been reported from other studies on invaded river systems where black-bass introduction has resulted in two distinct assemblages, one dominated by small native fishes and the other by large-bodied *Micropterus* spp.

(Godinho & Ferreira, 2000; MacRae & Jackson, 2001; van der Walt *et al.*, 2017). In *Micropterus* dominated assemblages, the number of other species present has also been shown to be directly related to habitat complexity and heterogeneity, providing refuge from predation (Tonn & Magnuson, 1982; Schlosser, 1987; MacRae & Jackson, 2001; Moyle *et al.*, 2003). Swartkops headwater tributaries are episodic and pools are predominantly fed by interstitial flows and isolated on the surface. During dry periods (>95% of the time) native species are confined, therefore, with predatory black basses in invaded pools. For fishes inhabiting episodic environments, their persistence relies on refugia not prone to desiccation (Magoulick & Kobza, 2003), but if these are occupied by invasive predatory fishes (as was the case on the Blindekloof River), the persistence of the species is severely compromised. This has been demonstrated in systems with seasonal rainfall fluctuations, such as the Cosumnes River, California, USA (Moyle *et al.*, 2003) and the Guadiana River Basin in Portugal (Collares-Pereira *et al.*, 1998), where seasonal droughts result in the extirpation of native fishes from isolated pools inhabited by black basses.

In situations where black-bass invasion results in the total extirpation or exclusion of native fishes from invaded zones, changes additional to loss of habitat become difficult to quantify. An above average rainfall year, however, provided an opportunity to corroborate stable state findings after *P. afer* recolonized black-bass invaded stream reaches, that were previously devoid of small native fishes. Over the monitoring period in the Blindekloof River, overall abundances of *P. afer* in pools inhabited by black basses were not significantly lower than in the non-invaded zone in the upper stream reaches, as well as in the mixed zone in the lower reaches. There were, however, size specific differences in temporal abundance between stream zones. The maintenance of *P. afer* in the invaded zone was most probably a result of dispersal into invaded stream reaches from non-invaded source populations. Similar meta-population dynamics, where the maintenance of native fishes in invaded zones is due to immigration from non-invaded source populations, have been observed for Canterbury galaxias *Galaxias vulgaris* Stokell 1949 in salmonid invaded streams in New Zealand (Woodford & McIntosh, 2010, 2011). Specifically, *G. vulgaris* only persisted in reaches invaded by *Oncorhynchus mykiss* (Walbaum 1792) and *Salmo trutta* L. 1758 when invader-free source populations were present (Woodford & McIntosh, 2011).

In the Blindekloof River, YOY *P. afer* increased in abundance over the sampling period, as this corresponded to the summer spawning season for this species (Cambray, 1994). Despite the extremely low abundance of adult *P. afer* in the invaded zone, YOY *P. afer* abundances did not differ significantly between invaded and non-invaded zones. This is contrary to findings by Godinho & Ferreira (2000), who found that in the Raia River in Portugal, juvenile abundance of native fishes was significantly lower in stream reaches invaded by non-native *M. salmoides* and pumpkinseed *Lepomis gibbosus* (L. 1758).

In the case of the Blindekloof River, the low abundance of black basses in invaded zones and the massive recruitment of YOY *P. afer* into the invaded zone probably resulted in a lag period between the influx and any noticeable changes in abundance due to predation over the relatively short duration of the study. In the mixed zone, similar increases in YOY *P. afer* abundances were observed, but they were significantly lower than in the invaded and non-invaded zones. Two possible causes for this are a decreased adult abundance in the mixed zone, resulting in poor recruitment (and thus lower downstream dispersal), or variable reproductive potential in stream habitats due

to physical or environmental factors making them less favourable in these respects than other stream reaches (Labbe & Fausch, 2000).

Adult *P. afer* abundance differed significantly between stream zones, with the low abundance in the invaded zone, intermediate abundance in the mixed zone and high abundance in the non-invaded zone. *Pseudobarbus afer* adult abundance also displayed temporal changes between zones. In the Blindekloof River, the invaded zone may therefore be acting as a filter or sink, resulting in reduced abundances of *P. afer* adults downstream, as this longitudinal decrease in abundance was not observed from the adjacent non-invaded Fernkloof and Waterkloof tributaries of the Swartkops River system (Ellender & Weyl, 2015). The influence of other non-native fishes (e.g., *T. sparrmanii* and *C. gariepinus*) on *P. afer* recruitment in the mixed zone can also not be discounted. There were, however, indications of adult dispersal from the upper to lower stream reaches.

In the non-invaded zone in the upper stream reaches there was a negative relationship between abundance and sample month and concomitantly the opposite was true for the mixed zone in the lower stream reaches, where there was an accumulation of *P. afer* adults over time. Despite the net movement of *P. afer* adults downstream, which continually flushed individuals through the invaded zone, by the end of the study adult *P. afer* were once again rare in the invaded zone and absent from the pool with the highest abundance of black basses. In pools where black basses were consistently recorded, *P. afer* abundance decreased rapidly to undetectable levels within 3 months. As a result of flows associated with a high rainfall month, adult *P. afer* were again observed in the invaded zone. Within the invaded zone, but at sites where black basses were not recorded, the opposite was true and *P. afer* abundance increased over time to >15 times that recorded at invaded sites.

Whether *P. afer* naturally displays source–sink population dynamics [where the upstream reaches act as a source and downstream reaches a demographic sink; (Pulliam & Danielson, 1991; Dias, 1996)] is not known. Alternatively, as has been documented for the Arkansas darter *Etheostoma cragini* Gilbert, 1885 in intermittent Colorado plains streams (Labbe & Fausch, 2000), *P. afer* may migrate downstream to use inundated habitats and migrate back upstream into refuge habitats as the stream dries up before the onset of the dry season. In the case of *E. cragini* the ability to disperse between refuge pools and habitats prone to seasonal drying was shown to be vital for rearing young of the year, which highlights the importance of connectivity at much finer spatial scales than previously considered (Schlosser, 1991; Labbe & Fausch, 2000; Fausch *et al.*, 2002). If *P. afer* display similar behaviour, the biological barrier created by the presence of black basses in certain stream segments may inhibit both dispersal and migration between important habitats. This seems most likely to affect adult *P. afer*, which exhibited the highest vulnerability to black-bass invasion.

A question arising is why native predators appear to have lesser effects than non-native predators. Separating the effect of non-native black basses and native predators (*A. mossambica*, *A. marmorata* and *S. capensis*) on the *P. afer* population is complex and some sort of natural predation pressure on *P. afer* has to be realistically assumed as all three species are reported to eat fish (Bruton *et al.*, 1987). *Anguilla marmorata* and *A. mossambica* were ubiquitous in headwater streams inhabited by *P. afer*. Having co-evolved in these headwater streams could explain why *P. afer* may

possibly also be less affected less by these native predators, than they are by non-native predators.

Naïveté by native fishes to introduced predators has been shown to result in large changes in native fish species (Whitlow *et al.*, 2003; Cox & Lima, 2006). From laboratory experiments Kadye & Booth (2014) alluded to inactivity and use of structure at night as evolved behaviour by *P. afer* to avoid native eel predation. During this study, nocturnal field observations of *P. afer* position in the water column provide additional insight into this. While observing *P. afer* behaviour during day and night snorkelling surveys conducted on the Blindekloof River, it was apparent that at night, *P. afer* did not use structure, but was predominantly suspended in the water column, well off the bottom of the stream while during the day, individuals were observed swimming in the water column and hiding in cracks and under rocks. At night, *Anguilla* spp. were observed hunting on the bottom of the stream, examining cracks and crevices while they systematically covered a pool. This suggests that the suspension in the water column is most likely an evolved predator-avoidance behaviour that might reduce predation rates by native eels. As predatory success of black basses has been shown to be negatively affected by habitat complexity (Savino & Stein, 1982; Alexander *et al.*, 2015) such behaviour increase the vulnerability of *P. afer* to predation.

In conclusion, evidence suggests that the invasion of black basses in the Blindekloof River may not disrupt YOY dispersal and recruitment, but rather their invasion of the middle stream reaches reduced adult abundance and hence also recruitment downstream. It remains to be seen whether *P. afer* can indefinitely maintain the current isolated metapopulations that characterize its distribution due to black-bass invasions. The only way to accurately determine long-term effects would be to establish whether these populations can maintain themselves, or whether they rely on inputs from others within the stream network. To effectively conserve small native stream fishes, a multi-level understanding of population processes at multiple scales (from reach to system) is needed to prioritize efforts for remediation and rehabilitation (Schlosser & Angermeier, 1995; Labbe & Fausch, 2000).

The authors would like to thank the National Research Foundation of South Africa (NRF), Rhodes University and the DST/NRF Centre of Excellence for Invasion Biology for financial support. C. Coppinger, G. Taylor, T. Murray, P. Gourley and T. Bellingan are thanked for assistance in the field. The research that this publication is based on was conducted according to protocols which comply with the ethical standards of the South African Institute for Aquatic Biodiversity. Permits for field surveys were issued by the Department of Economic Development and Environmental Affairs (DEDEA; permit no. CRO16/10CR, CRO 17/10) and Eastern Cape Parks Board. Eastern Cape Parks Board and the staff of the Groendal Wilderness Area are thanked for access and for their logistical assistance.

Supporting Information

Supporting Information may be found in the online version of this paper:

TABLE S1. The stream zones (mixed, invaded, non-invaded), 15 sites (including volume and water quality), distribution, mean abundance (fish 100 m⁻² ± S.E.) and relative abundance (%) of fishes by site, from site 1 in the lower reaches to site 15 in the upper reaches, that were monitored for 6 months (day and night snorkelling: November 2011–April 2012) on the Blindekloof River, a headwater tributary of the Swartkops

River within the Groendal Wilderness Area (*non-native species). Grey shading indicates presence for each particular species.

References

- Alexander, M. E., Kaiser, H., Weyl, O. L. F. & Dick, J. T. A. (2015). Habitat simplification increases the impact of a freshwater invasive fish. *Environmental Biology of Fishes* **98**, 477–486.
- Angermeier, P. L. & Winston, M. R. (1999). Characterizing fish community diversity across Virginia landscapes: prerequisite for conservation. *Ecological Applications* **9**, 335–349.
- ter Braak, C. J. F. (1995). Ordination. In *Data Analysis in Community and Landscape Ecology* (Jongman, R., ter Braak, C. & van Tongeren, O., eds), pp. 91–173. Wageningen: Pudoc.
- ter Braak, C. J. F. & Šmilauer, P. (1998). *CANOCO Reference Manual and User's Guide to CANOCO for Windows: Software for Canonical Community Ordination (Version 4)*. Biometris, Plant Research International: Wageningen. www.biometris-plant-research-international.software.informer.com/
- ter Braak, C. J. F. & Šmilauer, P. (2002). *CANOCO for Windows Version 4. 51*. Wageningen: Biometris, Plant Research International Available at www.biometris-plant-research-international.software.informer.com/
- Bruton, M. N., Bok, A. H. & Davies, M. T. T. (1987). Life history styles of diadromous fishes in inland waters of southern Africa. In *Common Strategies of Anadromous and Catadromous Fishes* (American Fisheries Society Symposium, Vol. 1 (Dadswell, M. J., Klauda, R. J., Moffitt, C. M. & Saunders, R. L., eds), pp. 104–121. Bethesda: American Fisheries Society.
- Cambray, J. A. (1994). The comparative reproductive styles of two closely related African minnows (*Pseudobarbus afer* and *P. asper*) inhabiting two different sections of the Gamtoos River system. *Environmental Biology of Fishes* **41**, 247–268.
- Clark, B. M., Impson, D. & Rall, J. (2009). Present status and historical changes in the fish fauna of the Berg River, South Africa. *Transactions of The Royal Society Of South Africa* **64**, 142–163.
- Collares-Pereira, M. J., Cowx, I. G., Rodrigues, J. A., Rogado, L. & da Costa, L. M. (1999). The status of *Anaocypris hispanica* in Portugal: problems of conserving a highly endangered Iberian fish. *Biological Conservation* **88**, 207–212.
- Collares-Pereira, M. J., Pires, A. M., Coelho, M. M. & Cowx, I. G. (1998). Towards a conservation strategy for *Anaocypris hispanica*, the most endangered non-migratory fish in Portuguese streams. In *Stocking and Introduction of Fish* (Cowx, I. G., ed), pp. 437–499. Oxford: Blackwell Science–Fishing News Books.
- Cox, J. G. & Lima, S. L. (2006). Naïveté and an aquatic–terrestrial dichotomy in the effects of introduced predators. *Evolution* **21**, 674–680.
- Cunjak, R. A. & Power, G. (1986). Winter habitat utilization by stream resident brook trout (*Salvelinus fontinalis*) and brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* **43**, 1970–1981.
- Dias, P. C. (1996). Sources and sinks in population biology. *Trends in Ecology & Evolution* **11**, 326–330.
- Ellender, B. R., Weyl, O. L. F. & Swartz, E. R. (2011). Invasion of a headwater stream by non-native fishes in the Swartkops River system, South Africa. *African Zoology* **46**, 39–46.
- Ellender, B. R. & Weyl, O. L. F. (2015). Resilience of imperilled headwater stream fish to an unpredictable high-magnitude flood. *Koedoe* **57**, 1–8.
- Ellender, B. R., Wasserman, R. J., Chakona, A., Skelton, P. H. & Weyl, O. L. F. (2017). A review of the biology and status of cape fold ecoregion freshwater fishes. *Aquatic Conservation: Marine and Freshwater Ecosystems* **27**, 867–879.
- Fausch, K. D., Torgersen, C. E., Baxter, C. V., Li, H. W. & Li, W. (2002). Landscapes to riverscapes: bridging the gap between research and conservation of stream fishes. *Bioscience* **52**, 483–498.

- Godinho, F. N. & Ferreira, M. T. (2000). Composition of endemic fish assemblages in relation to exotic species and river regulation in a temperate stream. *Biological Invasions* **2**, 231–244.
- Henriques, R., Sousa, V. & Coelho, M. M. (2010). Migration patterns counteract seasonal isolation of *Squalius torgalensis*, a critically endangered freshwater fish inhabiting a typical circum-Mediterranean small drainage. *Conservation Genetics* **11**, 1859–1870.
- Jackson, D. A., Peres-Neto, P. R. & Olden, J. D. (2001). What controls who is where in freshwater fish communities – the roles of biotic, abiotic and spatial factors. *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 157–170.
- Kulhanek, S. A., Ricciardi, A. & Leung, B. (2011). Is invasion history a useful tool for predicting the impacts of the world's worst aquatic invasive species? *Ecological Applications* **21**, 189–202.
- Kadye, W. T. & Booth, A. J. (2014). Alternative responses to predation in two headwater stream minnows is reflected in their contrasting diel activity patterns. *PloS One* **9**, p.e93666.
- Kimberg, P. K., Woodford, D. J., Roux, H. & Weyl, O. L. F. (2014). Species-specific impact of introduced largemouth bass *Micropterus salmoides* in the Groot Marico freshwater ecosystem priority area, South Africa. *African Journal of Aquatic Science* **39**, 451–458.
- Labbe, T. R. & Fausch, K. D. (2000). Dynamics of intermittent stream habitat regulate persistence of a threatened fish at multiple scales. *Ecological Applications* **10**, 1774–1791.
- MacRae, P. S. D. & Jackson, D. A. (2001). The influence of smallmouth bass (*Micropterus dolomieu*) predation and habitat complexity on the structure of littoral zone fish assemblages. *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 342–351.
- Magoulick, D. D. & Kobza, R. M. (2003). The role of refugia for fishes during drought: a review and synthesis. *Freshwater Biology* **48**, 1186–1198.
- Marr, S. M., Marchetti, M. P., Olden, J. D., García-Berthou, E., Morgan, D. L., Arismendi, I., Day, J. A., Griffiths, C. L. & Skelton, P. H. (2010). Freshwater fish introductions in mediterranean-climate regions: are there commonalities in the conservation problem? *Diversity and Distributions* **16**, 606–619.
- Meyer, J. L., Strayer, D. L., Wallace, J. B., Eggert, S. L., Helfman, G. S. & Leonard, N. E. (2007). The contribution of headwater streams to biodiversity in river networks. *Journal of the American Water Resources Association* **43**, 86–103.
- Moyle, P. B., Crain, P. K., Whitener, K. & Mount, J. F. (2003). Alien fishes in natural streams: fish distribution, assemblage structure and conservation in the Cosumnes River, California, USA. *Environmental Biology of Fishes* **68**, 143–162.
- Moyle, P. B. & Williams, J. E. (1990). Biodiversity loss in the temperate zone: decline of the native fish fauna of California. *Conservation Biology* **4**, 275–284.
- Parker, I. M., Simberloff, D., Lonsdale, W. M., Goodell, K., Wonham, M., Kareiva, P. M., Williamson, M. H., Von Holle, B., Moyle, P. B., Byers, J. E. & Goldwasser, L. (1999). Impact: toward a framework for understanding the ecological effects of invaders. *Biological Invasions* **1**, 3–19.
- Pulliam, H. R. & Danielson, B. J. (1991). Sources, sinks and habitat selection: a landscape perspective on population dynamics. *The American Naturalist* **137**, S50–S66.
- Ricciardi, A. & Atkinson, S. K. (2004). Distinctiveness magnifies the impact of biological invaders in aquatic ecosystems. *Ecology Letters* **7**, 781–784.
- RStudio Team (2015). *RStudio: Integrated Development for R*. Boston, MA: RStudio Inc. www.rstudio.com/.
- Savino, J. F. & Stein, R. A. (1982). Predator-prey interaction between largemouth bass and bluegills as influenced by simulated, submersed vegetation. *Transactions of the American Fisheries Society* **111**, 255–266.
- Schlosser, I. J. (1991). Fish ecology: a landscape perspective. *Bioscience* **41**, 704–712.
- Schlosser, I. J. & Angermeier, P. L. (1995). Spatial variation in demographic processes of lotic fishes: conceptual models, empirical evidence and implications for conservation. In *Evolution and the Aquatic Ecosystem: Defining Unique Units in Population Conservation* American Fisheries Society Symposium **17** (Nielsen, J. L., ed), pp. 392–401. Bethesda: American Fisheries Society.
- Schlosser, I. J. (1987). The role of predation in age- and size-related habitat use by stream fishes. *Ecology* **68**, 651–659.

- Smith, T. A. & Kraft, C. E. (2005). River fish assemblages in relation to landscape position and local habitat variables. *Transactions of the American Fisheries Society* **134**, 430–440.
- Strahler, A. N. (1957). Quantitative analysis of watershed geomorphology. *Transactions, American Geophysical Union* **38**, 913–920.
- Thurrow, R. & Schill, D. J. (1996). Comparison of day snorkeling, night snorkeling and electrofishing to estimate bull trout abundance and size structure in a second-order Idaho stream. *North American Journal of Fisheries Management* **16**, 314–323.
- Tonn, W. M. & Magnuson, J. T. (1982). Patterns in the species composition and richness of fish assemblages in northern Wisconsin lakes. *Ecology* **63**, 1149–1166.
- van der Walt, J. A., Weyl, O. L. F., Woodford, D. J. & Radloff, F. G. (2016). Spatial extent and consequences of black bass (*Micropterus* spp.) invasion in a cape floristic region river basin. *Aquatic Conservation: Marine and Freshwater Ecosystems* **26**, 736–748.
- van der Walt, K. A., Swartz, E. R., Woodford, D. J. & Weyl, O. L. F. (2017). Using genetics to prioritise headwater stream fish populations of the Marico barb, *Enteromius motebensis* Steindachner 1894, for conservation action. *Koedoe* **59**, 1–7.
- Weyl, O. L. F., Ellender, B. R., Woodford, D. J. & Jordaan, M. S. (2013). Native fish distributions and immediate impact of rotenone treatment on alien fish abundance in the Rondegat River, cape floristic region, South Africa. *African Journal of Aquatic Science* **38**, 201–209.
- Whitlow, W. L., Rice, N. A. & Sweeney, C. (2003). Native species vulnerability to introduced predators: testing an inducible defense and a refuge from predation. *Biological Invasions* **5**, 23–31.
- Woodford, D. J., Impson, N. D., Day, J. A. & Bills, I. R. (2005). The predatory impact of invasive alien smallmouth bass *Micropterus dolomieu* (Teleostei: Centrarchidae), on indigenous fishes in a cape floristic region mountain stream. *African Journal of Aquatic Science* **30**, 167–173.
- Woodford, D. J. & McIntosh, A. R. (2010). Evidence of source–sink metapopulations in a vulnerable native galaxiid fish driven by introduced trout. *Ecological Applications* **20**, 967–977.
- Woodford, D. J. & McIntosh, A. R. (2011). Location of demographic sources affects the distributions of a vulnerable native fish in invaded river networks. *Freshwater Biology* **56**, 311–324.