Specific spatio-temporal dynamics of absorptive fine roots in response to
neighbor species identity in a mixed beech-spruce forest

Zwetsloot, M. J., Goebel, M., Paya, A., Grams, T. E. E., & Bauerle, T. L.

This is a "Post-Print" accepted manuscript, which has been Published in "Tree
Physiology"

This version is distributed under a non-commercial no derivatives Creative Commons

([CC-BY-NC-ND](#)) user license, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited and not used for commercial purposes. Further, the restriction applies that if you remix, transform, or build upon the material, you may not distribute the modified material.

Please cite this publication as follows:

Zwetsloot, M. J., Goebel, M., Paya, A., Grams, T. E. E., & Bauerle, T. L. (2019). Specific spatio-temporal dynamics of absorptive fine roots in response to neighbor species identity in a mixed beech-spruce forest. *Tree Physiology*, 39(11), 1867-1879. <https://doi.org/10.1093/treephys/tpz086>

You can download the published version at:

<https://doi.org/10.1093/treephys/tpz086>

**Title: Specific spatio-temporal dynamics of absorptive fine roots in response to
neighbor species identity in a mixed beech-spruce forest**

Marie J Zwetsloot^{1,2}, Marc Goebel³, Alex Paya¹, Thorsten EE Grams⁴, and Taryn L
Bauerle^{1,5}

¹Cornell University, School of Integrative Plant Science, Ithaca, NY, USA

²Present address: Wageningen University, Soil Biology Group, Wageningen, The
Netherlands

³Cornell University, Department of Natural Resources, Ithaca, NY, USA

⁴Technical University of Munich, Ecophysiology of Plants, Freising, Germany

⁵Corresponding author (+1 607-254-4867, bauerle@cornell.edu)

Key words: belowground species interactions, absorptive fine roots, spatio-temporal
dynamics, root survivorship, root respiration, forest ecology

Main text: 6967 words

Abstract

Absorptive fine roots are an important driver of soil biogeochemical cycles. Yet, the spatio-temporal dynamics of those roots in the presence of neighboring species remains poorly understood. The aim of this study was to analyze shifts in absorptive fine-root traits in monoculture or mixtures of *Fagus sylvatica* [L.] and *Picea abies* [L.] Karst. We hypothesized that root competition would be higher under single-species than mixed-species interactions, leading to changes in 1) root survivorship, diameter, and respiration, and 2) spatio-temporal patterns of root growth and death. Using minirhizotron methods, we monitored the timing and location of absorptive fine-root growth and death at an experimental forest in southern Germany from 2011-2013. We also measured root respiration in the spring and fall seasons of 2012 and 2013. Our findings show that the absorptive fine roots of *F. sylvatica* had a 50% higher risk of root mortality and higher respiration rates in the single-species compared to the mixed-species zones. These results support our hypothesis that root competition is less intense for *F. sylvatica* in mixture versus monoculture. We were unable to find confirmation for the same hypothesis for *P. abies*. To analyze spatio-temporal patterns of absorptive fine-root production and mortality, we used a mixed-effects model considering root depth (space) and seasons (time) simultaneously. This analysis showed that *F. sylvatica* shifts root production towards shallower soil layers in mixed-species stands, besides significant seasonal fluctuations in root production depths for both species. Ultimately, the impact of neighbor species identity on root traits observed in this study have important implications for where, when and how fast root-facilitated carbon cycling takes place in single-species versus mixed-species forests. In addition, our study highlights the need for inclusion of absorptive fine-root spatio-temporal dynamics when examining belowground plant interactions and biogeochemical cycles.

Introduction

The spatio-temporal dynamics of fine-root growth are crucial for understanding biogeochemical cycles and belowground plant-plant interactions in a changing climate (Nadelhoffer & Raich 1992, Eissenstat et al. 2000, Gill & Jackson 2000, Bardgett et al. 2014). Fine roots and root-associated organisms are the primary sources of soil carbon (C) in forest ecosystems (Rasse et al. 2005, Clemmensen et al. 2013). Hence, the timing and location of fine-root production and mortality within the soil profile largely determines the distribution and cycling of belowground C resources. Even though root competition is a ubiquitous feature of plant communities and can significantly affect plant performance (Casper & Jackson 1997, Cahill & Casper 2000, Novoplansky 2009), it is unclear to what extent the species identity of neighboring plants influences fine-root traits and thereby soil C inputs in forests.

Recently, fine roots (< 2 mm in diameter) have been redefined into two distinct functional categories: (1) absorptive fine roots, the most distal, lowest-order roots with high resource uptake capacity, and higher incidence of mycorrhizal colonization, and (2) higher-order, transport roots with increased conductance rates based on their ontogeny and secondary structural development (McCormack et al. 2015). Absorptive, lower-order fine roots can include ephemeral modules, which are shorter-lived than transport, higher-order roots (Xia et al. 2010, Valenzuela-Estrade et al. 2008, Espeleta et al. 2009), and therefore constitute the faster turnover component of the fine-root system. The more dynamic character of absorptive fine roots with smaller diameters also allows them to be more responsive to changes in environmental factors (McCormack & Guo 2014) and interactions with neighboring tree species. Hence, their potentially greater role in belowground plant-plant interactions and impact on soil biogeochemical cycles made them the primary focus of this study.

The heterogeneous distribution of soil resources increases the risk of being outcompeted by neighboring plants and requires root foraging strategies over space and time (Casper & Jackson 1997, Farley & Fitter 1999, Ettema & Wardle 2002, Grams & Andersen 2007). Foraging strategies including timing of fine-root production, shifts in lifespan, morphology and physiology can all contribute to the intensity of root

interactions (Tilman 1988, Hutchings & de Kroon 1994, Hodge 2006, Volder et al. 2005, Schenk 1999). Root proliferation often coincides with periods of increased resource availability (Eissenstat & Caldwell 1988, Pregitzer et al. 1993). Studies have shown that plant species intrinsically vary the timing of root production to preempt a resource or spatially partition its uptake (Chesson 2000, Stratton et al. 2000, McKane et al. 2002). In addition, fine-root morphology can respond to resource availability. Because fine roots are responsible for resource acquisition and associations with beneficial soil microorganisms (Pregitzer et al. 1997, Pregitzer et al. 2002), the modification of fine-root diameter or length has critical consequences for resource uptake and plant growth (Eissenstat 1992). A reduction in diameter coupled with an increase in length results in a greater absorptive fine-root surface area without augmenting C and nitrogen costs for root production (Eissenstat 1991, Eissenstat et al. 2002, Ostonen et al. 2007). Even though these previous studies have thoroughly explored how root foraging strategies respond to shifts in resource availability and distribution, it is unknown whether fine-root traits show spatial and temporal adjustments to neighbor species identity.

The root foraging strategy that may have received the least attention in the context of species interactions is fine-root longevity or lifespan. Ultimately, resource exploitation does not solely depend on the level of fine-root production in a resource path but also on the lifespan of the roots occupying that space (Adams et al., 2013). Fine-root lifespan can be plastic in some species and change depending on resource availability (Aber et al. 1985, Burton et al. 2000, Adams et al., 2013). In a scenario where trees are competing for soil resources, individuals that maintain fine-roots within resource patches for longer periods of time can effectively preempt soil space and ensure a greater return on investment (Eissenstat & Yanai 1997, Eissenstat et al. 2000). Yet, the benefits of space occupation may be outweighed by the C costs of producing and maintaining fine roots. For example, fine-root respiration as a measure of maintenance costs is associated with soil temperature, fine-root activity and nitrogen content (Zogg et al., 1996; Volder et al., 2005) and, therefore, likely to be affected by seasonal variations in temperature and precipitation. Yet, it is unclear whether belowground neighbor species identity affects root lifespan across seasons.

The few experiments focusing on belowground plant-plant interactions in forest

ecosystems have shown that root foraging strategies differ between monotypic and mixed-species forest stands. Schmid & Kazda (2002) found that tree species shift their average rooting depth when grown in mixed-species arrangements compared to monocultures, essentially changing their niche within the soil profile. When discrete resource patches are unavailable due to a high proportion of root system overlap, roots of distinct tree species may also alter their timing of water uptake and level of root activity, allowing for the coexistence of neighbors with very similar niche requirements (Meinzer et al. 1999, Stratton et al. 2000). In addition, fine roots of *Fagus sylvatica* [L.] (European beech) showed greater surface area and specific root length when grown intermixed with *Picea abies* [L.] Karst (Norway spruce) than in monoculture (Bolte & Villanueva, 2006). Yet, other studies have reported no admixing effects on root morphology (Lei et al. 2012). In terms of fine-root lifespan, one study reported that fine-root longevity decreased for *F. sylvatica* and increased for *Fraxinus excelsior* (European ash) when grown together in the same area (Beyer et al. 2013). The authors attributed these opposite trends to asymmetric competition in favor of ash trees. However, because this is one of the few studies testing the effect of neighbor identity on fine-root lifespan, further experimentation is needed to test these trends.

At the ecosystem scale, previous studies have shown a positive relationship between species diversity and productivity (Hooper & Vitousek 1997, Hector 1999, Tilman 2001, Pretzsch et al. 2010, Liang et al. 2016). So far, most biodiversity-productivity research has focused on grassland ecosystems and the majority of these studies considered solely aboveground parameters (Meinen et al. 2009). Species differences in root traits facilitating niche complementary could explain increases in belowground root productivity observed in forests (Brassard et al. 2011). In the case where significant belowground niche overlap exists among different tree species, the pattern of increased belowground root productivity or “over-yielding” of species involved may not be evident (Meinen et al. 2009). From these previous findings on root productivity, the question arises to what extent the outcome of belowground species interactions can be explained by static species differences in root system traits or plastic root responses to neighboring trees.

We designed an experiment aimed at testing the effects of single-species versus

139 mixed-species root interactions on the timing and location of absorptive fine-root growth,
140 root survivorship and respiration of *F. sylvatica* and *P. abies* forest stands in southern
141 Germany over three years. We hypothesized that trees growing intermixed with a
142 different species (mixed-species) would experience less belowground competition in
143 comparison to trees growing in monoculture (single-species). We therefore predicted, for
144 both species, that absorptive fine roots produced within the single-species zones of *F.*
145 *sylvatica* or *P. abies* would have smaller average diameters (morphological adjustment),
146 lower cumulative root production, higher risk of mortality (lower survivorship), and
147 higher respiration rates (physiological adjustment) when compared to roots growing in
148 the mixed-species zone. Moreover, we expected to find evidence for niche
149 complementarity by identifying a significant difference in the timing and average depth
150 of absorptive fine-root production and mortality between the mixed-species and single-
151 species zones of the forest.

154 **Materials and methods**

156 *Experimental setup and field site conditions*

157 The study was performed across ten randomized plots containing 70-75-year-old, mature
158 *Picea abies* [L.] Karst (Norway spruce) and *Fagus sylvatica* [L.] (European beech) trees
159 located in the “Kranzberger Forst,” a research site near Freising, Germany (48°25’N,
160 11°39’E, 490 m a.s.l.). Total stand basal area was 52.0 m² ha⁻¹ with the stand area of *F.*
161 *sylvatica* amounting to 22.7 m² ha⁻¹ and that of *P. abies* to 29.3 m² ha⁻¹. Total stand
162 density was 659 trees ha⁻¹, of which 353 were *F. sylvatica* and 305 were *P. abies* (Rötzer
163 et al. 2017). Rectangular plots (~ 8 x 18 m) contained monotypic groupings of spruce
164 trees at one end, and at the opposite end, a monotypic grouping of beech trees (Pretzsch
165 et al. 2014). The central portion of each plot contained a “mixed” zone with a high degree
166 of interspecific root mixing (Mainiero et al. 2010, Häberle et al. 2012). For a map of the
167 research plots and tree distribution, please see Rötzer et al. (2017). To prevent cross-plot
168 root interactions and water flow between each of the ten experimental plots, plots were
169 trenched in May 2010 until a known clay-pan was reached (~ 1 meter), lined with low-

density poly-vinyl plastic, and then backfilled (Pretzsch et al. 2016).

Thirty-year averages (1970-2000) of mean precipitation and air temperature were 786 mm and 7.6°C respectively (Matyssek et al. 2007). During the 2011 – 2013 period, however, mean air temperature and annual precipitation were 8.0°C and 810 mm. Air temperature, relative humidity, and precipitation were collected continuously from the “Ecosystem Monitoring Plot Freising” (Bavarian State Institute of Forestry, LWF, Freising, Germany), 1 km south of the Kranzberger forest research site. Following Gaussen (1958) and Walter & Leith (1960), temporal occurrences of drought conditions, as well as, freeze periods can be observed at the site, i.e. the month of July in 2013, the winter seasons of 2012 and 2013 (Figure 1), respectively.

Soils at the site are haplic luvisols and provide sufficient nutrients for tree growth (Göttlein et al. 2012). Time domain reflectometers (TDR) used to monitor volumetric soil water content were installed in 2010 across all ten plots at two soil depths: 0-10 and 10-30 cm (Cole, 1977). Soil volumetric water content was measured weekly from April through November, and monthly from December through March (TDR-100, Campbell Scientific, North Logan, Utah). Averaged over three years, our measurements showed that average soil water content in mixed ($22.6 \pm 0.1\%$) and beech-dominated soils ($21.9 \pm 0.1\%$) in the 0-10 cm depth was higher compared to spruce-dominated soils ($20.3\% \pm 0.1$, supplementary figure 1). Deeper soil layers (10-30 cm) also had higher average soil water content in mixed ($28.5 \pm 0.1\%$) and beech-dominated stands ($28.1 \pm 0.1\%$) compared to spruce-dominated stands ($26.2 \pm 0.1\%$). During the month of August (2011-2013), the monotypic beech and spruce as well as the mixed-species zones reached equally low average moisture levels of $18.9 \pm 0.5\%$ at 0-10 cm and $23.6 \pm 0.5\%$ at 10-30 cm soil depth.

Root Observation

In 2010, clear acrylic minirhizotron tubes (70 cm long, 6 cm outside diameter) were installed at an angle of 60° from the horizontal reaching a vertical depth of 50 cm. The vertical depth was chosen based on previous research results at Kranzberger forest, which reported that > 90% of beech and spruce roots grew between 0-50cm (Häberle *et al.* 2012). Each plot contained six minirhizotron tubes: two in each single-species zone, from

hereon called “*F. sylvatica*” and “*P. abies*” zone, and two within the beech-spruce zone, from hereon called “mixed-species” zone. Each tube was located a minimum distance of one meter from the plot boundaries, and in the case of the mixed-species zones, tubes were installed equidistant from both species. Average distances from the minirhizotron tubes to the trees were significantly longer in the mixed-species zone (5.4 m for *F. sylvatica* and 5.5 m for *P. abies*) than in the single-species zone (2.8 m for *F. sylvatica* and 2.9 m for *P. abies*, $P < 0.001$). However, minirhizotron tube distance to trees did not have a significant effect on cumulative fine root production within these tubes ($P = 0.83$, supplementary figure 2). Moreover, we did not find a significant difference in tube distances to trees between the two species ($P = 0.67$). Before installation, minirhizotron tubes were capped at the base with plastic plugs and lined with silicon caulk to reduce water infiltration. Tubes, when not in use, were covered with large plastic caps to prevent aboveground water infiltration and light penetration.

From the 18th of May 2011 until the 29th of August 2013, 44 continuous images (width = 18 mm, height = 15 mm) were taken across the length of each tube using a minirhizotron camera system (BTC100X Camera, Bartz Technology, Carpinteria, California), bi-weekly during the growing season and monthly during the winter months. All images were analyzed for absorptive fine-root production, death, survivorship, and diameter using *WinRHIZO Tron MF* (Regent Inc., Quebec, Canada). To be certain of the absorptive fine-root characteristic, we solely focused on first-order roots (root tips). If a first-order root grew into a second-order root, this was marked as such and excluded from data analysis. The date of root birth was calculated as the mid-point between the appearance of a new root tip and the preceding image session. Similarly, the death date of a root was calculated as the mid-point between the disappearance or death of a root (black, shriveled) and the preceding image session (Comas et al. 2005). Roots that transected more than one observation window were only counted once. The number of neighboring roots was defined as the number of root tips present in the proximity (within the 18 mm by 15 mm image) of the root at the time of death. Through visual inspection of images in the single-species zones of *F. sylvatica* and *P. abies*, we determined three root specific differences by appearance: (1) the epidermal coloration (spruce: dark brown, beech: reddish) (2) root tip branching patterns (spruce: alternate branching, beech:

herringbone and often opposite branching), and (3) root tip ending (spruce: round, beech: sharp). These three indicators were used to identify species identity of roots in the mixed-species zones.

Root Respiration

To view, trace, and sample roots of known species and age (Bouma et al. 2001), three in-ground plywood boxes (root box: 40 x 60 x 30 cm, H x W x D) containing two acetate windows were installed in eight out of the ten research plots; one in each single-species zone, and one in the mixed-species zone. Viewing windows of the root boxes, when not in use, were insulated year-round with polystyrene shutters, and a lid was screwed into place to prevent micro-climatic effects and exclude light penetration.

Root respiration was measured in the spring and fall of 2012 (May 22 and October 6), and 2013 (May 18 and October 7). Using the root boxes, absorptive fine-root branches containing first- and second-order roots were traced along acetate windows using colored markers for a period of three weeks to ensure that fine-root material was of a known species and age ($5 < x < 22$ days old). Pioneer roots, defined as first-order roots exhibiting little or no branching, lower incidence of mycorrhizal colonization, and larger average diameter (Zadworny & Eissenstat 2011) were excluded in this experiment. Highlighted roots were then excised from behind the acetate windows using a razor blade, transferred to an onsite climate-controlled hut, and placed into a cuvette containing oxygenated buffer (10 mM MES, 1mM CaSO₄, and 5.5 μ M K₂HPO₄, pH = 5.8; Volder et al. 2005) within an oxygen electrode system maintained at 21 ± 0.5 °C (Oxygraph, Hansatech, King's Lynn, UK). Oxygen consumption was measured over a period of 15 minutes. After 15 minutes, roots were removed from the cuvette, placed in a paper envelope, and dried at 60°C.

Data Analysis

All statistical analyses were performed in R (R Core Team, 2018). We used the emmeans (Lenth 2019), survival and coxme (Thernau & Grambsch 2000), and lme4 (Bates et al. 2015) packages for Tukey Honest Significance Difference (HSD) tests, survival analyses and mixed-effects linear models.

263 To determine the effect of species interaction zones (single-species *F. sylvatica*,
264 single-species *P. abies*, and mixed-species) on belowground root productivity measured
265 as cumulative root tip production, we used a linear model and Tukey HSD tests. Root tip
266 production of each species interaction zone was calculated by averaging the production of
267 first-order roots over the two tubes in the mixed-species and single-species zone of each
268 replicate plot and expressed as the total number of tips produced per square meter
269 minirhizotron viewing window. To analyze root tip production of the species interaction
270 zones by 10 cm soil depth increments and seasons, we used the same approach but log-
271 transformed seasonal root tip production to satisfy the normality and equal variance of
272 the residuals assumptions. Seasons were defined as winter (December 01-February 28),
273 spring (March 01-May 31), summer (June 01- August 31), and fall (September 01-
274 November 30). Root tip mortality patterns were analyzed using the same methods.

275 To analyze shifts in absorptive fine-root production and mortality patterns of *F.*
276 *sylvatica* and *P. abies* over space and time, we again analyzed first-order roots only and
277 took two different modeling approaches. First, we used two separate linear models to test
278 the effect of neighbor identity on the proportion of fine-root tips by 10 cm depth
279 increments or season separately. The proportion was calculated by dividing the number
280 of root tips in a specific temporal or spatial cohort of each year by the cumulative annual
281 root tip production or death over the 50 cm soil profile. Secondly, we developed a spatio-
282 temporal mixed-effects model where both depth (responding variable) and seasons
283 (independent variable) were included simultaneously. In addition, we included species
284 neighbor identity, year, and an interaction term for neighbor species identity and season
285 as independent variables in the model. Plot was included as a random effect.

286 For the survivorship analysis of absorptive fine roots, we made use of Cox
287 proportional hazards regressions (Cox, 1972). Neighbor species identity, root diameter,
288 number of neighboring root tips and root depth were included as covariates and plot as a
289 random effect. To satisfy Cox's proportional hazard assumptions, (1) we stratified the
290 fine root data in the model by season and production year, and (2) we categorized the
291 independent variables root diameter (0-0.2 mm, 0.2-0.4 mm, > 0.4 mm) and root depth
292 (0-10 cm, 10-30 cm, 30-50 cm) into cohorts.

293 Lastly, we used a linear model and Tukey HSD tests to determine the effect of

species, neighbor species identity and season on fine-root respiration. This was the only exception where second-order roots were also assessed. To analyze trends in first-order root diameter, we utilized a mixed-effects model with plot as random effect.

Results

Spatio-temporal patterns of absorptive fine-root production and mortality

Cumulative root tip production was greater in the single-species zone of *F. sylvatica* than the single-species zone of *P. abies* during 2011-2013 (Figure 2a, $P < 0.05$). Yet, cumulative root tip production in the mixed-species zone was not significantly different from the single-species zones for either species, implying no evidence for over-yielding under mixed-species interactions. In mixture, *F. sylvatica* produced a greater number of root tips ($46.6 \cdot 10^2$ tips m^{-2} window) than *P. abies* ($11.8 \cdot 10^2$ tips m^{-2} window). These trends in root tip production under single- and mixed-species growing conditions persisted at different spatial and temporal scales: at 20-30 cm, 30-40 cm and 40-50 cm soil depth (Figure 2b), and across seasons (Supplementary figure 3). The ratio of *F. sylvatica* to *P. abies* root tips increased with soil depth.

Cumulative root tip deaths in the single- and mixed-species zones during 2011-2013 showed the same pattern as production (Supplementary figure 4). At greater soil depth, the single-species zone of *F. sylvatica* had a greater number of root tip deaths than the single-species zone of *P. abies* (Supplementary figure 5a). Root tip death was greater in the single-species zone of *F. sylvatica* than *P. abies* in the summer and fall, yet such differences were not observed in winter and spring (Supplementary figure 5b). Whether we analyzed the total amount of dead tips during 2011-2013 or partitioned the data by season or depth, we did not find a difference in root tip death between the single- and mixed-species zones except for the single *P. abies* zone and mixture in the fall.

Using the proportion of root tips produced over 10 cm soil depth levels or seasons by year as a responding variable, we found differences in the spatial root production patterns of *F. sylvatica* and *P. abies* under single- and mixed-species growing conditions (Figure 3). Namely, while *P. abies* had a greater proportion of absorptive fine roots

growing in the upper 10 cm of the soil when grown in mixture with *F. sylvatica* than by itself, *F. sylvatica* had a greater proportion of fine roots growing at 10-20 cm soil depth in the mixed-species arrangement ($P < 0.05$). In general, *F. sylvatica* showed an even distribution of absorptive fine roots across depth levels, while *P. abies* produced a large proportion of root tips in the upper 10 cm of the soil that decreased with soil depth. Temporal adjustments in the form of increased absorptive fine-root production, was only observed in the summer for the single-species zone of *F. sylvatica* relative to mixed-species growing conditions. Overall, root tip production varied seasonally across years for both *F. sylvatica* and *P. abies*, with most roots produced in the summer followed by spring and fall. Fine-root production was lowest in the winter for both species ($P < 0.05$). Also, for both species, root tip production was the highest in 2011 ($P < 0.05$). Root tip production of *P. abies* was similar in 2012 and 2013, while *F. sylvatica* produced more fine roots in 2012 than 2013 ($P < 0.05$). It is important to note that we stopped measuring root tip production in September 2013, and more absorptive fine roots may have been produced in the fall and winter of that year.

When including soil depth (space as dependent variable) and season (time as one of the independent variables) in the same mixed-effects model, we found that absorptive fine roots of *F. sylvatica* were produced deeper in the soil (25.5 ± 3.1 cm) when grown with itself than in mixed-species stands (21.1 ± 3.1 cm, $P < 0.001$) (Figure 4a), while *P. abies* root tip production depth was not affected by the identity of neighboring species ($P = 0.85$). Likewise, we did not detect a significant interaction between single- versus mixed-species zones and production season, meaning that *F. sylvatica* and *P. abies* did not show temporal adjustments in root tip growth dynamics in response to neighbor species identity ($P = 0.13$ and $P = 0.17$, respectively).

The depth of root tip production differed by season for both species but was more pronounced for *F. sylvatica* than *P. abies*. Of the total population of *F. sylvatica* roots produced, fall roots were produced significantly deeper in the soil compared to other seasons ($P < 0.05$). Root tip production in spring took place at the shallowest depth but was not significantly different from the average production depth in winter. As the seasons progressed from spring-summer-fall, root tips were produced at continuously deeper depths ($P < 0.05$). For *P. abies*, absorptive fine roots in summer and fall were

produced deeper than spring roots ($P < 0.05$) but the production depth of winter roots was not significantly different from the rooting depth of the other seasons. For both species, root tip production in 2011 was shallower than the other years. For *F. sylvatica*, rooting depth was also shallower in 2012 than 2013. Overall, average root production depth of *P. abies* (15.2 ± 2.6 cm) was shallower than that of *F. sylvatica* (23.6 ± 2.6 cm).

Average depth of root tip mortality in the single-species and mixed-species stands followed the same trend as fine-root production (Figure 4b) with two exceptions. First, we did not detect a significant effect of species interaction zones for *F. sylvatica* in the spring. Second, we found a significant interaction between year and species interaction zones for *P. abies* ($P < 0.001$), which showed a difference in the depth of root tip mortality between the single-species (17.8 ± 2.6 cm) and mixed-species (7.1 ± 3.2 cm) arrangements in 2012 but not in 2011 or 2013. Although, the lack of significant difference in 2013 may be due to the absence of fall data. Interestingly, in the case of *F. sylvatica*, seasonal patterns of root tip mortality were the inverse of production with root tip death taking place in the shallowest soil layers in the summer and fall, and at greater depth in spring followed by winter ($P < 0.05$). In addition, root tip death of *F. sylvatica* took place at a progressively deeper depth each year ($P < 0.05$). For *P. abies*, season did not influence the depth of fine-root mortality ($P = 0.25$).

Survivorship of absorptive fine roots

The root tips of *F. sylvatica* lived longer in mixed-species growing conditions (Table 1, supplementary figure 6a). The Cox proportional hazard model estimated that beech absorptive fine roots in the single-species zone have a 54% higher risk of mortality compared to roots in mixed-species zones ($P < 0.0001$), with a median lifespan of 174 days versus 440 days, respectively. In contrast to *F. sylvatica*, *P. abies* did not show a significant difference in absorptive fine-root survivorship under mixed- or single-species conditions (Table 2, supplementary figure 6b). On the other hand, soil depth significantly influenced the survivorship of both *F. sylvatica* and *P. abies* root tips. The depth effect was stronger for *F. sylvatica* ($P < 0.0001$) than for *P. abies* ($P = 0.025$) at 10-30 cm in comparison to the reference 0-10 cm surface soil layer (Table 1, 2). Absorptive fine-root lifespan increased for both species in deeper soil layers (Supplementary figure 7). The

median root lifespan of *F. sylvatica* was 147 (0-10 cm), 220 (10-30 cm) and 400 (30-50 cm) days. Yet, for *P. abies*, only the deeper roots at 30-50 cm had a significantly higher lifespan (318 days) than the upper layers which showed similar lifespans (234 days, 0-10 cm and 190 days, 10-30 cm). A smaller resolution of 10 cm depth increments also resulted in no depth effect differences. *F. sylvatica* absorptive fine roots with a diameter > 0.2 mm lived longer than the roots with a diameter < 0.2 mm (Table 1, $P < 0.0001$). In the case of *P. abies*, however, only root tips with a diameter > 0.4 mm had longer lifespans than the 0-0.2 mm diameter cohort (Table 2, $P = 0.035$). The number of neighboring root tips present at the time of death did not have a significant effect on root survivorship for either species (Table 1, 2).

Morphological and physiological absorptive fine-root plasticity to neighbor species identity

Neighbor identity significantly influenced absorptive fine-root respiration for *F. sylvatica* in spring of 2012 and 2013 and for *P. abies* in the spring of 2012 (Figure 5) with a significant third-order interaction among year, season and species interaction zones. In all these cases, absorptive fine-root respiration rates were higher in the single-species zone than in the mixed-species zone. For both species, season significantly influenced root respiration ($P < 0.0001$). In 2013, respiration was higher in the fall than in the spring for both species under single-species and mixed-species conditions ($P < 0.05$). In 2012, *F. sylvatica* roots in the mixed-species zone respired more in the fall than in the spring ($P < 0.05$), while the trend was reverse for single-species roots of *F. sylvatica* ($P < 0.05$). For *P. abies*, we did not find a significant seasonal difference in root respiration under mixed-species conditions in 2012 ($P = 0.99$). Yet, the single-species absorptive fine roots of *P. abies* showed higher respiration in the spring than in the fall of 2012 ($P < 0.05$). Overall, absorptive fine roots of *P. abies* had higher respiration rates in 2013 than in 2012 ($P < 0.05$), but root respiration of *F. sylvatica* did not vary between years ($P = 0.51$). Absorptive fine-root respiration did not differ between species ($P = 0.28$).

Root tips of *F. sylvatica* were smaller in diameter than those of *P. abies* ($P < 0.05$). Absorptive fine roots of *F. sylvatica* in the mixed-species zone were also slightly

larger than roots in the single-species zone (0.28 vs. 0.27 mm, $P < 0.05$), while the opposite was true for spruce (0.36 vs. 0.40 mm, $P < 0.05$). Also, the season in which root tips were produced had a significant effect on the absorptive fine-root diameter of both species. Absorptive fine roots of *F. sylvatica* had a smaller diameter in the summer and fall than in the spring and winter ($P < 0.05$). The diameter of *P. abies* absorptive fine roots was the largest in the spring, followed by summer and lastly by fall and winter ($P < 0.05$).

Discussion

Plastic responses in absorptive fine-root traits to neighbor species identity

This study analyzed absorptive fine-root traits and spatio-temporal dynamics of root production and mortality of *F. sylvatica* and *P. abies* in single-species and mixed-species zones in an experimental forest in southern Germany. As we predicted, the species identity of surrounding trees significantly affected the absorptive fine-root lifespan of *F. sylvatica* (Table 1, supplementary figure 6), respiration in spring (Figure 5), and diameter of both *F. sylvatica* and *P. abies*. We used these changes in absorptive fine-root traits as an indicator of the intensity of root competition in the single-species and mixed-species interaction zones. Contrary to our hypothesis that root competition is higher under single-species compared to mixed-species interactions for both *F. sylvatica* and *P. abies*, the directional shifts in fine-root traits observed in this study suggest that this is only the case for *F. sylvatica*. Our prediction on niche differentiation over space (soil depth) and time (seasons) was partly met and more dynamic than expected (Figure 3, 4).

To our knowledge, this study provides the first evaluation of absorptive fine-root survivorship in mature single-species and mixed-species forest stands at the field scale. We found that the risk of root tip mortality for *F. sylvatica* was 54% higher when grown by itself than when intermixed with *P. abies*. The higher rates of absorptive fine-root respiration of *F. sylvatica* in the single-species zone in comparison to the mixed-species zone in the spring of 2012 and 2013 line up well with the survivorship results. Namely, cost-benefit models predict that roots with higher respiration rates – indicating greater C costs - should have shorter lifespans (Eissenstat 1992, Eissenstat & Yanai 1997,

Eissenstat et al. 2000). These trends in root longevity and respiration may also be connected to changes in fine-root diameter. While previous research has demonstrated that fine roots with larger diameters tend to have lower respiration rates (Pregitzer et al. 1998) and longer root lifespans (Wells & Eissenstat 2001), these studies assessed different root orders with a wider range of diameters, soil depths and functions than covered in our analysis. The absorptive fine roots of *F. sylvatica* indeed had larger diameters in the mixed-species than single-species zone of the forest. Yet, this difference only amounted to 0.01 mm. Because few studies have analyzed the impact of changes in fine-root diameter at this scale, it is difficult to conclude whether such a minor change in diameter contributes to explaining the respiration and survivorship patterns observed in this study.

The absorptive fine roots of *P. abies* showed a less plastic response in survivorship and respiration (Table 2, Figure 5), but a more plastic response in diameter to neighbor species identity than *F. sylvatica*. Absorptive fine-root diameters of *P. abies* were 0.04 mm smaller under mixed-species than single-species interactions, the opposite response to that of *F. sylvatica*. A smaller fine-root diameter associated with higher specific root length is a common morphological adjustment to low soil resources (Farley & Fitter 1999, Hodge 2004). Hence, the species differences in root diameter, together with the observed lifespan patterns of *F. sylvatica* suggest that the soil environment under mixed-species interactions provided less competition and/or more resources to the absorptive fine roots of *F. sylvatica* but more competition and/or fewer resources to the fine roots of *P. abies* in comparison to their single-species environments.

Our conclusion that the mixed-species zone is more favorable for *F. sylvatica* but less favorable for *P. abies* than their respective single-species zones is supported by the observed trends in root space occupation. Namely, the total number of root tips produced substantially decreased from single *F. sylvatica* (beech roots), to mixed-species (beech + spruce roots), to single *P. abies* (spruce roots) (Figure 2). Though not significant, this trend suggests that mixed-species interactions provide an environment with fewer roots (reduced competition) for *F. sylvatica* yet increase root competition for *P. abies*. Another potential factor contributing to shifting absorptive fine-root traits between the single-species and mixed-species zone is soil resource availability. Yet, the lower soil moisture

content in the single *P. abies* zone in comparison to the mixture (Supplementary figure 1) goes against our conclusion that the mixed-species zone is a less favorable environment for *P. abies* root growth or suggests that competition for soil water resources was not driving shifts in absorptive fine-root traits. Previous studies at the “Kranzberger Forst” site identified higher nutrient concentrations in the topsoil in single *P. abies* than single *F. sylvatica* stands, which were primarily attributed to species-specific litter chemistry (Göttlein et al. 2012). This study, however, did not address soil nutrient content in mixed stands making it difficult to confirm the role of soil nutrient shifts. Lastly, while our study focused on fine-root growth dynamics at the forest stand or plot scale, it would be relevant to consider neighborhood conditions and specifically surrounding tree density effects on fine-root traits when shifting to the individual tree scale. Overall, our findings stress the complexity of belowground plant-plant interactions and that the outcome of root competition may be asymmetric (Montgomery et al. 2010, Rewald & Leuschner 2009). Future studies are needed to confirm the role of root space occupation, soil resource availability and tree density as drivers of shifts in fine-root morphology and physiology during belowground species interactions.

Spatio-temporal niche differentiation and fine-root production patterns under single-species and mixed-species root interactions

To assess patterns of spatial and temporal niche differentiation between the single- and mixed-species interaction zones of *F. sylvatica* and *P. abies*, we utilized two different analysis methods : 1) we evaluated the spatial or temporal proportion of root tips as individual response variables in two models, and 2) we developed one spatio-temporal model to analyze root depth as a response variable with season as the independent variable. Both modeling approaches gave similar results for spatial and temporal adjustments in absorptive fine-root production to neighboring tree species. First, we found that the single-species and mixed-species forest stands influence the spatial root distribution of *F. sylvatica*. Namely, *F. sylvatica* produced a greater proportion of root tips at 10-20 cm soil depth overall (Figure 3a, b) and had a shallower root production depth (Figure 4a) in the mixed-species zone in comparison to the single-species zone. The insignificant effect of neighbor species identity on the root production depth of *P.*

abies in the spatio-temporal model (Figure 4a) may be explained by its overall shallow root depth, which makes it more difficult to identify the significantly higher proportion of absorptive fine roots produced in the 0-10 cm soil layer in the mixed-species zone (Figure 3a, b). Schmid & Kazda (2002) also reported a shift in root production towards shallower soil layers for *P. abies* when grown in mixture. *F. sylvatica*'s root production on the other hand took place at deeper locations in the mixed zone. The fact that our study only focused on the absorptive fine roots – and not the entire root system – may explain these opposing trends. Secondly, neither model provided evidence for seasonal shifts in absorptive fine-root production between the single-species and mixed-species interaction zones apart from *F. sylvatica* producing higher proportion of root tips in summer under single-species than mixed-species composition in the one-dimensional seasonal model (Figure 3c, d).

The spatial root adjustments of *F. sylvatica* and *P. abies* to neighbor identity may go beyond mechanisms of resource availability (Schenk 2006, de Kroon 2007). Recently, root exudates and rhizosphere microbes have been recognized as important drivers of belowground plant-plant interactions (Mommer et al. 2016). Previous studies have shown distinct root exudate profiles (Zwetsloot et al. 2018) and fine-root fungal compositions (Nickel et al. 2018) of spruce and beech tree species. These relations between species-specific non-resource cues and spatial niche differentiation within the soil profile motivate further investigation on this topic.

We recognize that our second spatio-temporal model is a less common approach to analyzing absorptive fine-root dynamics. Yet, in addition to confirming the patterns observed by the first models that evaluated root dynamics separately over space and time, the second model highlights a spatio-temporal root tip production pattern that cannot be identified otherwise. Specifically, we found a significant effect of season on *F. sylvatica* and *P. abies* rooting depth (Figure 4a). This seasonal trend in root production depth has been observed in other studies (Lyr & Hoffman, 1967; Hendrick & Pregitzer, 1992) and could be explained by seasonal fluctuations in soil temperature and resource availability (Reich 1980) which are strongly associated to soil temperature (Teskey & Hinckley 1981; Kuhns et al. 1985; Tierney et al. 2003). For example, after bare soil exposure to low temperatures, shallow soil depths are the first to reach adequate temperatures for root

growth in spring. In addition, studies have observed that deep root production is related to soil water availability (Hendrick & Pregitzer, 1996). In our study, absorptive fine-root production in deeper soil layers may be a result of the lower soil moisture levels near the soil surface during the summer and fall months (Supplementary figure 1). Lastly, the location of absorptive fine-root mortality in the soil profile for *F. sylvatica* was the inverse of its production patterns through time (Figure 3b). This makes sense when we consider seasonal resource availability, where root tips are produced in areas of high nutrient or water content (shallow layers in spring, deep layers in fall) and senesce under unfavorable resource conditions (deep layers in spring, shallow layers in fall).

Belowground root productivity and implications for root-facilitated C cycling

Surprisingly, absorptive fine-root production in the mixed-species zone was not different from the single-species zone of *F. sylvatica* or *P. abies* (Figure 2a). This goes against the hypotheses and results of previous publications (Tilman et al. 1997, Brassard et al. 2008, Mommer et al. 2012) where species mixtures led to greater productivity than monocultures. The fact that we, and other studies (Meinen et al. 2008), did not find evidence of belowground over-yielding under mixed-species composition suggests that transitioning from a single-species setting to a mixture of species is not a guarantee for increased belowground root productivity. These contrasting outcomes of belowground plant-plant interactions could be explained by the extent of niche differentiation of the species involved in these different experiments. In our study, the absorptive fine roots of both species showed substantial niche overlap in the shallow soil layers. In light of competition and accessibility to soil resources, the overlap of absorptive fine roots may explain why belowground root productivity was not higher under mixed-species zones than in the single-species zones. However, Mommer et al. (2012) showed that mixtures of grass species had shallower rooting depths but also greater belowground root productivity than monocultures. An alternative hypothesis by Callaway & Walker (1997) states that competition is more prevalent under high resource availability, while facilitation is more common when resources are low. As in other similar studies, our conclusions are inherently limited to select species combinations under specific environmental conditions. Therefore, additional studies are needed to explore which mechanisms are

driving the outcomes of root-root interactions across ecosystems under fluctuating soil resources.

Cumulative absorptive fine-root production (Figure 2a) and mortality (Supplementary figure 4) in the species interaction zones showed similar trends. Yet, when we analyzed root tip mortality and production dynamics over soil depth and seasons (Figure 4), we found distinct patterns. With absorptive fine roots functioning as a primary source of soil C in forest ecosystems (Rasse et al. 2005, Clemmensen et al. 2013) and driving seasonal trends in microbial community composition and soil extracellular enzyme activity (Kaiser et al. 2010), the spatio-temporal patterns of absorptive fine-root mortality have important implications for where and when absorptive fine roots prime soil microbial processes and provide C inputs to soil organic matter pools. In contrast with the single-species zones of *P. abies*, where the depth of absorptive fine-root mortality did not fluctuate, or remained shallow in mixed zones, the mortality of absorptive fine roots of *F. sylvatica* shifted C fluxes from deeper soil layers in the spring to shallower depths in the summer and fall. When we consider total absorptive fine-root deaths in each season (Supplementary figure 5), our results imply that root C supply in the single- and mixed-species zone of *F. sylvatica* fluctuates across soil depths throughout the year with stronger pulses in the summer and fall. Few studies have attempted to analyze these seasonal and depth-related fine-root patterns, or they show contrasting results (Hendrick & Pregitzer 1996), making it challenging to extract general trends across species and experimental sites. We encourage future studies to make use of the spatio-temporal analyses presented in this paper to obtain a more complete understanding of root-facilitated C cycling in forest ecosystems.

Conclusion

In summary, this study analyzed the absorptive fine-root responses of *F. sylvatica* and *P. abies* in single- and mixed-species forest stands over space and time. Our findings have important implications for plant-plant interactions and biogeochemical cycles at the ecosystem scale. Namely, the observation that the absorptive fine roots of *F. sylvatica* show differences in longevity between the mixture and monoculture, and those of *P.*

604 *abies* do not, demonstrates that the plasticity of root lifespan in response to neighboring
605 plants is species-specific. Moreover, these results suggest slower turnover of *F. sylvatica*
606 absorptive fine roots and therefore root-facilitated C cycling in mixed-species forests. In
607 addition, we did not detect an increase in belowground root productivity in mixed-
608 relative to single-species stands. Our results confirm the findings from previous studies
609 that increasing tree species diversity is not a guarantee for higher root productivity.
610 Lastly, our spatio-temporal analyses of root growth dynamics indicate that root carbon
611 fluxes to the soil fluctuate within the soil depth profile throughout the year.

612 On a mechanistic level, future studies are necessary to disentangle what factors
613 drive shifts in absorptive fine-root dynamics in response to neighbor species identity. We
614 propose that soil resource availability, rhizodeposition, litter chemistry and species-
615 specific soil microbial communities likely contribute to spatial root niche differentiation
616 and changes in absorptive fine-root lifespan, respiration, and diameter between single-
617 species and mixed-species forest stands.

620 **Acknowledgements**

621 We are thankful for the invaluable technical support provided by Christof Seidler,
622 Thomas Feuerbach, Peter Kuba, Josef Heckmair, Annika Huber, and Roman Meier. We
623 also thank Karl-Heinz Häberle for his help during the entirety of this project and Stephen
624 Parry from Cornell Statistical Consulting Unit for his advice on analyzing spatio-
625 temporal patterns of root production and mortality.

628 **Conflict of Interest**

629 The authors declare no conflict of interest.

632 **Funding**

633 Andrew W. Mellon Student research grant and Einaudi funding to Alex Paya. Grants to
634 Thorsten Grams by the German Research Foundation (DFG, GR1881/5-1) and by the

Bavarian State Ministries of the Environment and Consumer Protection as well as Food,
Agriculture and Forestry.

References

- Aber JD, Melillo JM, Nadelhoffer KJ, Mcclaugherty CA, Pastor J (1985) Fine root turn-
over in forest ecosystems in relation to quantity and form of nitrogen availability–
a comparison of 2 methods. *Oecologia* 66: 317–321.
- Adams TS, McCormack ML, Eissenstat DM (2013) Foraging strategies in trees of
different root morphology: the role of root lifespan. *Tree Physiol* 33: 940-948.
- Bardgett RD, Mommer L, De Vries FT (2014) Going underground: root traits as drivers
of ecosystem processes. *Trends Ecol Evol* 29:692–699.
- Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software* 67: 1-48
- Beyer F, Hertel D, Jung K, Fender A-C, Leuschner C. 2013. Competition effects on fine
root survival of *Fagus sylvatica* and *Fraxinus excelsior*. *For Ecol Manage* 302:
14–22.
- Bolte A, Villanueva I (2006) Interspecific competition impacts on the morphology and
distribution of fine roots in European beech (*Fagus sylvatica* L.) and Norway
spruce (*Picea abies* (L.) Karst.). *Eur J For Res* 125:15-26.
- Bouma TJ, Yanai RD, Elkin AD, Hartmond U, Flores-Alva DE, Eissenstat DM (2001)
Estimating age-dependent costs and benefits of roots with contrasting life span:
comparing apples and oranges. *New Phytol* 150: 685-695.
- Brassard BW, Chen HYH, Bergeron Y, Paré D (2011) Differences in fine root
productivity between mixed- and single-species stands. *Funct Ecol* 25: 238–246.
- Burton AJ, Pregitzer KS, Hendrick RL (2000) Relationships between fine root dynamics
and nitrogen availability in Michigan northern hardwood forests. *Oecologia* 128:
389-399
- Callaway RM, Walker LR (1997) Competition and facilitation: a synthetic approach to
interactions in plant communities. *Ecology* 78: 1958-1965.

665 Cahill JF, Casper BB (2000) Investigating the relationship between neighbor root
 666 biomass and belowground competition: field evidence for symmetric competition
 667 belowground. *Oikos* 90:311–320.
 668 Casper BB, Jackson RB (1997) Plant competition underground. *Annu Rev Ecol Syst*
 669 28:545–570.
 670 Clemmensen KE, Bahr A, Ovaskainen O, Dahlberg A, Ekblad A, Wallander H, Stenlid
 671 J, Finaly RD, Wardle DA, Lindahl BD (2013) Roots and Associated Fungi Drive
 672 Long-Term Carbon Sequestration in Boreal Forest. *Science* 339: 1615-1618.
 673 Chesson P (2000) Mechanisms of maintenance of species diversity. *Annu Rev Ecol Syst*
 674 31: 343–36.
 675 Cole RH (1977) Time domain reflectometry. *Annu Rev Phys Chem* 28: 283-300.
 676 Comas LH, Eissenstat D (2004) Linking fine root traits to maximum potential growth rate
 677 among 11 mature temperate tree species. *Funct Ecol* 18: 388-397.
 678 Comas LH, Andersen LJ, Dunst RM, Lakso A, Eissenstat DM (2005) Canopy and
 679 environmental control of root dynamics in a long-term study of Concord grape.
 680 *New Phytol* 167: 829–840.
 681 Cox DR (1972) Regression models and life-tables. *J R Stat Soc Series B* 34: 187-202.
 682 Eissenstat DM, Caldwell MM (1988) Seasonal timing of root growth in favorable
 683 microsites. *Ecology*: 870-873.
 684 Eissenstat DM (1991) On the relationship between specific root length and the rate of
 685 root proliferation: a field study using citrus rootstocks. *New Phytol* 118: 63-68.
 686 Eissenstat DM (1992) Costs and benefits of constructing roots of small diameter. *J Plant*
 687 *Nutr* 15: 763-782.
 688 Eissenstat DM, CE Wells, RD Yanai, Whitbeck JL (2000) Building fine roots in a
 689 changing environment: implications for root longevity. *New Phytol* 147: 33-42.
 690 Eissenstat DM, Yanai RD (1997) The ecology of root lifespan. *Adv Ecol Res* 27: 1-60.
 691 Espeleta JF, West JB, Donovan LA (2009) Tree species fine-root demography parallels
 692 habitat specialization across a sandhill soil resource gradient. *Ecology* 90:1773–
 693 1787.
 694 Ettema CH, Wardele DA (2002) Spatial soil ecology. *Trends Ecol Evol* 17: 177-183.
 695 Farley RA, Fitter AH (1999) The responses of seven co-occurring woodland herbaceous

696 perennials to localizes nutrient-rich patches. *J Ecol* 87: 849-859.
 697 Gausсен H, de Phillips A (1958) La limite eumédi- terranéenne et les contrées de
 698 transition. Sous com- mission de coordination de la FAO des quations forestières
 699 méditerranéennes. FAO, Rome, Italy.
 700 Gill RA, Jackson RB (2000) Global patterns of root turnover for terrestrial ecosystems.
 701 *New Phytol* 147:13–31.
 702 Göttlein A, Baumgarten M, Dieler J (2012) Site Conditions and Tree-Internal Nutrient
 703 Partitioning in Mature European Beech and Norway Spruce at the Kranzberger
 704 Forst. In: Matyssek R, Schnyder H, Oßwald W, Ernst D, Munch JC, Pretzsch H
 705 (eds) *Growth and Defence in Plants: Resource Allocation at Multiple Scales*.
 706 Springer Berlin Heidelberg, Berlin, Heidelberg, pp 193–211.
 707 Grams TEE, Andersen CP (2007) Competition for resources in trees: Physiological
 708 versus morphological plasticity. In: Esser K, Lüttge U, Beyschlag W, Murata J
 709 (eds) *Progress in Botany*, vol 68. Springer-Verlag Berlin Heidelberg, pp 356-381.
 710 Häberle K-H, Weigt R, Nikolova PS, Reiter IM, Cermak J, Wieser G (2012) Case study:
 711 Kranzberger Forst”: Growth and Defense in European beech (*Fagus sylvatica*
 712 (L.)) and Norway spruce (*Picea abies* (L.) Karst.). In: Matyssek R, Schnyder H,
 713 Oßwald W, Ernst D, Munch JC, Pretzsch H (eds) *Growth and Defence in Plants:*
 714 *Resource Allocation at Multiple Scales*. Springer-Verlag Berlin Heidelberg, pp
 715 243-271.
 716 Hector A (1999) Plant Diversity and Productivity Experiments in European Grasslands.
 717 *Science* 286: 1123–1127.
 718 Hendrik RL, Pregitzer KS (1992) The demography of fine roots in a Northern hardwood
 719 forest. *Ecology* 73: 1094-1104.
 720 Hendrick RL, Pregitzer KS (1996) Temporal and depth-related patterns of fine root
 721 dynamics in northern hardwood forest. *J Ecol* 84: 167-176.
 722 Hodge A (2004) The plastic plant: root responses to heterogeneous supplies of nutrients.
 723 *New Phytol* 162: 9–24.
 724 Hodge A (2006) Plastic plants and patchy soils. *J Exp Bot* 57: 401-411.
 725 Hooper DU, Vitousek PM (1997) The effects of plant composition and diversity on
 726 ecosystem processes. *Science* 277: 1302-1305.

- Hutchings MJ, de Kroon H (1994) Foraging in plants: the role of morphological plasticity in resource acquisition. *Adv Ecol Res* 25: 159-238
- Kaiser C, Koranda M, Kitzler B, Fuchslueger L, Schnecker J, Schweiger P, Rasche F, Zechmeister-Boltenstern S, Sessitsch A, Richter A (2010) Belowground carbon allocation by trees drives seasonal patterns of extracellular enzyme activities by altering microbial community composition in beech forest soil. *New Phytol* 187: 843-858.
- de Kroon H (2007) How do roots interact? *Science* 318: 1562-1563.
- Kuhns MR, Garret HE, Teskey RO, Hinckley TM (1985) Root growth of black walnut trees related to soil temperatures, soil water potential and leaf water potential. *Forest Sci* 31: 617-629.
- Lei P, Scherer-Lorenzen M, Bauhus J. 2012. Belowground facilitation and competition in young tree species mixtures. *For Ecol Manage* 265: 191–200.
- Lenth R (2019) emmeans: Estimated Marginal Means, aka Least-Square Means. R package version 1.2.3: <https://CRAN.R-project.org/package=emmeans>
- Lyr H, Hoffman G (1967) Growth rates and growth periodicity of tree roots. *International Review of Forestry Research Volume 2*. Academic Press Inc, New York, pp 181-236.
- Liang J, Crowther TW, Picard N, Wiser S, Zhou M, Alberti G, Schulze E-D, McGuire AD, Bozzato F, Pretzsch H, et al. 2016. Positive biodiversity-productivity relationship predominant in global forests. *Science* 354: aaf8957.
- Mainiero R, Kazda M, Schmid I (2010) Fine root dynamics in 60-year-old stands of *Fagus sylvatica* and *Picea abies* growing on haplic luvisol soil. *Eur J For Res* 129: 1001-1009.
- Matyssek R, Bahnweg G, Ceulemans P, Fabien P, Grill D, Hanke DE, Kraigher H, Oßwald W, Rennenberg H, Sandermann H, Tausz M, Wieser G (2007) Synopsis of the CASIROZ Case Study: Carbon Sink Strength of *Fagus sylvatica* L. in a Changing Environment- Experimental Risk Assessment of Mitigation by Chronic Ozone Impact. *Plant Biol* 9: 163-180.
- McCormak ML, Guo D (2014) Impact of environmental factors on fine root lifespan. *Front Plant Sci* 5: 1-11.

758 McCormack ML, Dickie IA, Eissenstat DM, Fahey TJ, Fernandez CW, Guo D,
 759 Helmissaari H-S, Hobbie EA, Iversen CM, Jackson RB, Leppälammi-Kujansuu J,
 760 Norby RJ, Phillips RP, Pregitzer KS, Pritchard SG, Rewald B, Zadworny M
 761 (2015) Redefining fine roots improves understanding of below-ground
 762 contributions to terrestrial biosphere processes. *New Phytol* 207:505–518.

763 McKane RB, Johnson LC, Shaver GR, Nadelhoffer KJ, Rastetter EB, Fry B, Giblin AE,
 764 Kielland K, Kwiatkowski BL, Laundre JA, Murray G (2002) Resource-based
 765 niches provide a basis for plant species diversity and dominance in arctic tundra.
 766 *Nature* 415: 68-71.

767 Meinen C, Hertel D, Leuschner C (2009) Biomass and morphology of fine roots in
 768 temperate broad-leaved forests differing in tree species diversity: is there evidence
 769 of below-ground overyielding? *Oecologia* 161: 99–111.

770 Meinzer FD, Andrade JL, Goldstein G, Holbrook NM, Cavelier J, Wright SJ (1999)
 771 Partitioning of soil water among canopy trees in a seasonally dry tropical forest.
 772 *Oecologia* 121: 293-301.

773 Mommer L, Kirkegaard J, van Ruijven J (2016) Root-root interactions: towards a
 774 rhizosphere framework. *Trends Plant Sci* 21: 209-217.

775 Mommer L, van Ruijven J, de Caluwe H, Smit-Tiekstra AE, Wagemaker CAM, Ouborg
 776 NJ, Boegemann GM, van der Weerden GM, Berendse F, de Kroon H (2010)
 777 Unveiling below-ground species abundance in a biodiversity experiment: a test of
 778 vertical niche differentiation among grassland species. *J Ecol* 98: 1117-1127.

779 Montgomery RA, Reich P, Palik BJ (2010) Untangling positive and negative biotic
 780 interactions: views from above and below ground in a forest ecosystem. *Ecology*
 781 91: 3641-3655.

782 Nadelhoffer KJ, Raich JW (1992) Fine Root Production Estimates and Belowground
 783 Carbon Allocation in Forest Ecosystems. *Ecology* 73:1139–1147.

784 Novoplansky A (2009) Picking battles wisely: plant behaviour under competition. *Plant*
 785 *Cell Environ* 32:726–741.

786 Nickel UT, Weigl F, Kerner R, Schaefer C, Kallenbach C, Munch J, Pritsch K (2018)
 787 Quantitative losses vs. qualitative stability of ectomycorrhizal community
 788 responses to 3 years of experimental summer drought in a beech-spruce forest.

789 Glob Chang Biol 24: e560-e576.
 790 Ostonen I, Püttsepp Ü, Biel C, Alberton O, Bakker MR, Löhmus K, Metcalfe D,
 791 Olsthoorn AFM, Pronk A, Vanguelova E, Weigh M, Brunner I (2007) Specific
 792 root length as an indicator of environmental change. Plant Biosyst 141: 426-442.
 793 Pregitzer KS, Hendrick RL, Fogel R (1993) The demography of fine roots in response to
 794 patches of water and nitrogen. New Phytol 125: 575-580.
 795 Pregitzer KS, Kubiske ME, Yu CK, Hendrick RL (1997) Relationships among root
 796 branch order, carbon, and nitrogen in four temperate species. Oecologia 111: 302-
 797 308.
 798 Pregitzer KS, Laskowski MJ, Burton AJ, Lessard VC, Zak DR (1998) Variation in sugar
 799 maple root respiration with root diameter and soil depth. Tree Physiol 18: 665-
 800 670.
 801 Pregitzer KS, DeForest JL, Burton AJ, Allen MF, Ruess RW, Hendrick RL (2002) Fine
 802 root architecture of nine North American trees. Ecol Monogr 72: 293-309.
 803 Pretzsch H, Block J, Dieler J, Dong PH, Kohnle U, Nagel J, Spellmann H, Zingg A
 804 (2010) Comparison between the productivity of pure and mixed stands of Norway
 805 spruce and European beech along an ecological gradient. Ann For Sci 67: 712-
 806 712.
 807 Pretzsch H, Rötzer T, Matyssek R, Grams TEE, Häberle K-H, Pritsch K, Kerner R,
 808 Munch J-C (2014) Mixed Norway spruce (*Picea abies* L. Karst) and European
 809 beech (*Fagus sylvatica* L.) stands under drought: from reaction pattern to
 810 mechanism. Trees 28:1305–1321.
 811 Pretzsch H, Bauerle T, Häberle K, Matyssek R, Schütze G, Rötzer T (2016) Tree
 812 diameter growth after root trenching in a mature mixed stand of Norway spruce
 813 (*Picea abies* L. Karst) and European beech (*Fagus sylvatica* L.). Trees 30: 1761-
 814 1773.
 815 Rasse DP, Rumpel C, Dignac M (2005) Is soil carbon mostly root carbon? Mechanisms
 816 for specificic stabilization. Plant Soil 269: 341-356.
 817 R Core Team (2018) R: A language and environment for statistical computing. R
 818 Foundation for Statistical Computing, Vienna, Austria.
 819 Reich PB (1980) Period root and shoot growth in oak. Forest Sci 26: 590-597.

820 Rewald B, Leuschner B (2009) Belowground competition in a broad-leaved temperate
 821 mixed forest: patterns analysis and experiments in a four-species stand. *Eur J*
 822 *Forest Res* 128: 387-398.

823 Rötzer T, Biber P, Moser A, Schäfer C, Pretzsch H (2017) Stem and root diameter
 824 growth of European beech and Norway spruce under extreme drought. *Forest*
 825 *Ecology and Management* 406: 184–195.

826 Schenk HJ, Callaway RM, Mahall BE (1999) Spatial root segregation: are plants
 827 territorial? *Adv Ecol Res* 28:145–180.

828 Schenk HJ (2006) Root competition: beyond resource depletion. *J Ecol* 94: 725-739.

829 Schmid I, Kazda M (2002) Root distribution of Norway spruce in monospecific and
 830 mixed stands on different soils. *Forest Ecology and Management* 159:37–47.

831 Stratton LC, Goldstein G, Meinzer FC (2000) Temporal and spatial partitioning of water
 832 resources among eight woody species in a Hawaiian dry forest. *Oecologia* 124:
 833 309-317.

834 Teskey RO, Hinckley TM (1981) Influence of temperature and water potential on root
 835 growth of white oak. *Physiol Plant* 52: 363-369.

836 Thernau TM, Grambsch PM (eds) (2000). *Modeling Survival Data: Extending the Cox*
 837 *Model*. Springer, New York.

838 Tierney C, Fahey TJ, Groffman PM, Hardy JP, Fitzhugh RD, Driscoll CT, Yavitt JB
 839 (2003) *Glob Chane Biol* 9: 760-679.

840 Tilman D (ed) (1988) *Plant strategies and the dynamics and structure of plant*
 841 *communities* (No. 26). Princeton University Press, Princeton, NJ.

842 Tilman D (2001) Diversity and Productivity in a Long-Term Grassland Experiment.
 843 *Science* 294: 843–845.

844 Tilman D, Lehman CL, Thomson KT (1997) Plant diversity and ecosystem productivity:
 845 theoretical considerations. *Proc Natl Acad Sci* 94: 1857-1861.

846 Valenzuela-Estrada LR, Vera-Caraballo V, Ruth LE, Eissenstat DM (2008) Root
 847 anatomy, morphology, and longevity among root orders in *Vaccinium*
 848 *corymbosum* (Ericaceae). *Am J Bot* 95:1506–1514.

849 Volder A, Smart DR, Bloom AJ, Eissenstat DM (2005) Rapid decline in nitrate uptake
 850 and respiration with age in fine lateral roots of grape: implications for root

851 efficiency and competitive effectiveness. *New Phytol* 165: 493-502.
 852 Walter H, Lieth H (1960) *Klimadiagramm Weltatlas*. Gustav Fischer, Jena, Germany.
 853 Wells CE, Eissenstat DM (2001) Marked differences in survivorship among apple roots
 854 of different diameters. *Ecology* 8: 882-892.
 855 Xia M, Guo D, Pregitzer KS (2010) Ephemeral root modules in *Fraxinus mandshurica*.
 856 *New Phytol* 188:1065–1074.
 857 Zadworny M, Eissenstat DM (2011) Contrasting the morphology, anatomy and fungal
 858 colonization of new pioneer and fibrous roots. *New Phytol* 190: 213-221.
 859 Zogg GP, Zak DR, Burton AJ, Pregitzer KS. 1996. Fine root respiration in northern
 860 hardwood forests in relation to temperature and nitrogen availability. *Tree Physiol*
 861 16: 719–725.
 862 Zwetsloot MJ, Kessler A, Bauerle T (2018) Phenolic root exudate and tissue compounds
 863 vary widely among temperate forest tree species and have contrasting effects on
 864 soil microbial respiration. *New Phytol* 218: 530-541.
 865
 866

Figures

Figure 1. Average air temperature (point-line) and monthly total precipitation (bars) during 2011-2013 are in fixed proportion, 10°C corresponding to a precipitation of 20 mm, allowing the characterization of seasonal environmental conditions, i.e. drought or freeze periods (Gaussen 1958, Walter & Lieth 1960). During 2011-2013, mean air temperature and annual precipitation were 8.0 °C and 810 mm. Seasons are indicated by greyscale and defined as winter (Dec, Jan, Feb), spring (Mar, Apr, May), summer (June July, Aug), and fall (Sept, Oct, Nov).

Figure 2. Cumulative root tip production by (a) zones and (b) depth levels and zones of *F. sylvatica*, *P. abies*, and mixed-species zone (roots of both species combined together). Means and standard errors are shown ($N = 10$). Contrasting letters indicate significant differences among zones within depth level ($P < 0.05$) as determined with the Tukey Honest Significant Difference test. The area below the white line indicates the amount of roots belonging to *P. abies* and the area above the white line indicates the amount of roots belonging to *F. sylvatica* in the mixed-species zone.

Figure 3. Relative abundance of root tip production by 10 cm soil depth levels for (a) *F. sylvatica* and (b) *P. abies*, and by season for (c) *F. sylvatica* and (d) *P. abies* averaged over the years 2011-2013 under mixed-species (black) and single-species (grey) growing conditions. Means and standard errors are shown ($N = 10$). Contrasting uppercase letters indicate significant differences between depth levels or season averaged over single- and mixed-species interaction zones ($P < 0.05$) and contrasting lowercase letters indicate significant differences between mixed-species and single-species growing conditions ($P < 0.05$) as determined with the Tukey Honest Significant Difference test. Dashed line indicates 25% of cumulative root production in panel a and b, and 50% of cumulative root production in panel c and d.

Figure 4. Average depth $\pm$ standard error of (a) root tip production ($N = 4855$) and (b) mortality ($N = 3694$) across seasons and year for root tips in the single-species zone of *F. sylvatica* and *P. abies* as well as both species in the mixed-species zone. Means are indicated by the line graph and standard errors by grey-tinted area on each side of the line. Seasons are defined as: winter (Dec, Jan, Feb), spring (Mar, Apr, May), summer (June July, Aug), and fall (Sept, Oct, Nov).

Figure 5. Respiration of absorptive fine roots (first- and second-order roots, 5 < x < 22 days old) of (a) *F. sylvatica* and (b) *P. abies* growing under single-species (grey) and mixed-species (black) conditions in the spring (22nd of May 2012, 18th of May 2013) and fall (6th of October, 7th of October) season. Means and standard errors are shown ($N = 56$ for *F. sylvatica* and $N = 56$ for *P. abies* in 2012, $N = 48$ for *F. sylvatica* and $N = 46$ for *P. abies* in 2013). Contrasting letters indicate significant differences between seasons and single- and mixed-species interaction zones within each year as determined by Tukey Honest Significant Difference tests ($P < 0.05$).

Tables

Table 1. Cox-proportional hazard model output for the mortality risk of root tips of *F. sylvatica*. Coefficient estimates, standard error (SE), Wald Chi-Square statistic (χ^2), hazard risk ratio and p-values are shown. For the categorical variables included in the model, their reference level is in parentheses. Significant p-values ($P < 0.05$) are in bold. A hazard ratio < 1 means a decrease in mortality risk, while a hazard ratio > 1 signifies an increase in mortality risk. Neighbors are defined as number of neighboring root tips of *F. sylvatica* and *P. abies* in 18 mm by 15 mm image at time of death.

Variable	Coefficient estimate	SE	χ^2	<i>P</i>	Hazard ratio
Interaction (Mixed-species) Single-species	0.43	0.05	8.62	< 0.0001	1.54
Diameter (0-0.2 mm)					
0.2-0.4 mm	-0.58	0.05	-12.46	< 0.0001	0.56
> 0.4 mm	-1.40	0.17	-8.06	< 0.0001	0.25
Depth (0-10 cm)					
10-30 cm	-0.31	0.06	-4.87	< 0.0001	0.73
30-50 cm	-0.91	0.07	-12.34	< 0.0001	0.40
Neighbors	-0.001	0.001	-1.02	0.31	1.00

Table 2. Cox-proportional hazard model output for the mortality risk of root tips of *P. abies*. Coefficient estimates, standard error (SE), Wald Chi-Square statistic (χ^2), hazard risk ratio and p-values are shown. For the categorical variables included in the model, their reference level is in parentheses. Significant p-values ($P < 0.05$) are in bold. A hazard ratio < 1 means a decrease in mortality risk, while a hazard ratio > 1 signifies an increase in mortality risk. Neighbors are defined as number of neighboring root tips of *F. sylvatica* and *P. abies* in 18 mm by 15 mm image at time of death.

Variable	Coefficient estimate	SE	χ^2	<i>P</i>	Hazard ratio
Interaction (Mixed-species) Single-species	-0.02	0.11	-0.13	0.89	0.99
Diameter (0-0.2 mm)					
0.2-0.4 mm	-0.20	0.29	-0.66	0.51	0.82
> 0.4 mm	-0.65	0.31	-2.11	0.04	0.52
Depth (0-10 cm)					
10-30 cm	-0.23	0.10	-2.24	0.03	0.79
30-50 cm	-1.06	0.19	-5.52	< 0.0001	0.34
Neighbors	0.001	0.001	0.46	0.65	1.00