

LIST OF REFERENCES

- Abrams, P.A., 1988. How should resources be counted? *Theoretical Population Biology*, 33 (2), 226-242.
- Abrams, P.A., 1990. The effects of adaptive behavior on the type-2 functional response. *Ecology*, 71 (3), 877-885.
- Abrams, P.A., 1995. Implications of dynamically variable traits for identifying, classifying, and measuring direct and indirect effects in ecological communities. *American Naturalist*, 146 (1), 112-134.
- Abrams, P.A., 2000. The impact of habitat selection on the spatial heterogeneity of resources in varying environments. *Ecology*, 81 (10), 2902-2913.
- Aerts, B.A., Esselink, P. and Helder, G.J.F., 1996. Habitat selection and diet composition of greylag geese *Anser anser* and barnacle geese *Branta leucopsis* during fall and spring staging in relation to management in the tidal marshes of the Dollard. *Zeitschrift für Ökologie und Naturschutz*, 5 (2), 65-75.
- Albon, S.D. and Langvatn, R., 1992. Plant phenology and the benefits of migration in a temperate ungulate. *Oikos*, 65 (3), 502-513.
- Allen, A.W., Cook, J.G. and Armbruster, M.J., 1984. *Habitat suitability index models: pronghorn*. US Fish and Wildlife Service no. FWS/OBS-82/10.65. [http://el.ercd.usace.army.mil/emrrp/emris/EMRIS_PDF/Pronghorn.pdf]
- Alonso, D., Bartumeus, F. and Catalan, J., 2002. Mutual interference between predators can give rise to Turing spatial patterns. *Ecology*, 83 (1), 28-34.
- Anderson, A., 1984. The extinction of moa in southern New Zealand. In: Martin, P.S. and Klein, R.G. eds. *Quaternary extinctions: a prehistoric revolution* University of Arizona Press, Tucson, 728-740.
- Anderson, J.A., 1995. *An introduction to neural networks*. MIT Press, Cambridge.
- Andresen, H., Bakker, J.P., Brongers, M., et al., 1990. Long-term changes of salt marsh communities by cattle grazing. *Vegetatio*, 89 (2), 137-148.
- ARC, 1980. *The nutrient requirements of ruminant livestock: technical review by an Agricultural Research Council working party*. Agricultural Research Council, Commonwealth Agricultural Bureaux, Slough.
- Arditi, R. and Dacorogna, B., 1988. Optimal foraging on arbitrary food distributions and the definition of habitat patches. *American Naturalist*, 131 (6), 837-846.
- Arnold, G.W., 1985. Regulation of forage intake. In: Hudson, R.J. and White, R.G. eds. *Bioenergetics of wild herbivores*. CRC Press, Boca Raton, 81-101.
- Arnold, W., Ruf, T., Reimoser, S., et al., 2004. Nocturnal hypometabolism as an overwintering strategy of red deer (*Cervus elaphus*). *American Journal of Physiology*, 286 (1), R174-R181.
- Arsenault, R. and Owen-Smith, N., 2002. Facilitation versus competition in grazing herbivore assemblages. *Oikos*, 97 (3), 313-318.
- Asner, G.P., 1998. Biophysical and biochemical sources of variability in canopy reflectance. *Remote Sensing of Environment*, 64 (3), 234-253.
- Astrom, M., Lunberg, P. and Danell, K., 1990. Partial prey consumption by browsers: trees as patches. *Journal of Animal Ecology*, 59 (1), 287-300.
- Atkinson, M.D., Jervis, A.P. and Sangha, R.S., 1997. Discrimination between *Betula pendula*, *Betula pubescens*, and their hybrids using near-infrared reflectance spectroscopy. *Canadian Journal of Forest Research*, 27 (11), 1896-1900.
- Atwood, S.B., Provenza, F.D., Wiedmeier, R.D., et al., 2001a. Changes in preferences of gestating heifers fed untreated or ammoniated straw in different flavours. *Journal of Animal Science*, 79 (12), 3027-3033.
- Atwood, S.B., Provenza, F.D., Wiedmeier, R.D., et al., 2001b. Influence of free-choice vs. mixed-ration diets on food intake and performance of fattening calves. *Journal of Animal Science*, 79 (12), 3034-3040.
- Augustin, N.H., Muggleston, M.A. and Buckland, S.T., 1996. An autologistic model for the spatial distribution of wildlife. *Journal of Applied Ecology*, 33 (2), 339-347.

- Bailey, D.W., 1995. Daily selection of feeding areas by cattle in homogeneous and heterogeneous environments. *Applied Animal Behaviour Science*, 45 (3/4), 183-199.
- Bailey, D.W., 2004. Management strategies for optimal grazing distribution and use of arid rangelands. *Journal of Animal Science*, 82 (E-Suppl.), E147-E153.
- Bailey, D.W., Gross, J.E., Laca, E.A., et al., 1996. Mechanisms that result in large herbivore grazing distribution patterns. *Journal of Range Management*, 49 (5), 386-400.
- Bailey, D.W., Howery, L.D. and Boss, D.L., 2000. Effects of social facilitation for locating feeding sites by cattle in an eight-arm radial maze. *Applied Animal Behaviour Science*, 68 (2), 93-105.
- Bailey, D.W., Keil, M.R. and Rittenhouse, L.R., 2004. Research observation: daily movement patterns of hill climbing and bottom dwelling cows. *Journal of Range Management*, 57 (1), 20-28.
- Bailey, D.W., Kress, D.D., Anderson, D.C., et al., 2001a. Evaluation of F1 crosses from Angus, Charolais, Salers, Piedmontese, Tarentaise and Hereford sires. V. Grazing distribution patterns. *Proceedings of the Western Section American Society of Animal Science*, 52, 110-113.
- Bailey, D.W., Kress, D.D., Anderson, D.C., et al., 2001b. Relationship between terrain use and performance of beef cows grazing foothill rangeland. *Journal of Animal Science*, 79 (7), 1883-1891.
- Bailey, D.W., Rittenhouse, L.R., Hart, R.H., et al., 1989a. Characteristics of spatial memory in cattle. *Applied Animal Behaviour Science*, 23 (4), 331-340.
- Bailey, D.W., Rittenhouse, L.R., Hart, R.H., et al., 1989b. Association of relative food availabilities and locations by cattle. *Journal of Range Management*, 42 (6), 480-482.
- Bailey, D.W. and Sims, P.L., 1998. Association of food quality and locations by cattle. *Journal of Range Management*, 51 (1), 2-8.
- Bailey, D.W. and VanWagoner, H., 2004. Can we improve livestock distribution with selection? In: *Abstracts of the International Society for Range Management Annual Meeting*. Salt Lake City.
- Bailey, D.W., Walker, J.W. and Rittenhouse, L.R., 1990. Sequential analysis of cattle location: day-to-day movement patterns. *Applied Animal Behaviour Science*, 25 (1/2), 137-148.
- Bailey, D.W. and Welling, G.R., 1999. Modification of cattle grazing distribution with dehydrated molasses supplement. *Journal of Range Management*, 52 (6), 575-582.
- Bailey, D.W., Welling, G.R. and Miller, E.T., 2001c. Cattle use of foothills rangeland near dehydrated molasses supplement. *Journal of Range Management*, 54 (4), 338-347.
- Bailey, R.G., 1996. *Ecosystem geography*. Springer, New York.
- Baker, D.L. and Hobbs, N.T., 1987. Strategies of digestion: digestive efficiency and retention time of forage diets in montane ungulates. *Canadian Journal of Zoology*, 65 (8), 1978-1984.
- Bakker, J.P., 1989. *Nature management by grazing and cutting: on the ecological significance of grazing and cutting regimes applied to restore former species-rich grassland communities in the Netherlands*. Kluwer Academic Publishers, Dordrecht. Geobotany no. 14.
- Ball, J.P., Danell, K. and Sunesson, P., 2000. Response of a herbivore community to increased food quality and quantity: an experiment with nitrogen fertilizer in a boreal forest. *Journal of Applied Ecology*, 37 (2), 247-255.
- Baret, F., Jacquemoud, S., Guyot, G., et al., 1992. Modeled analysis of the biophysical nature of spectral shifts and comparison with information content of broad bands. *Remote Sensing of Environment*, 41 (2/3), 133-142.
- Barry, T.N., Manley, T.R. and Duncan, S.J., 1986. The role of condensed tannins in the nutritional value of *Lotus pedunculatus* for sheep. *British Journal of Nutrition*, 55 (1), 123-137.
- Bashaw, M.J., Tarou, L.R., Maki, T.S., et al., 2001. A survey assessment of variables related to stereotypy in captive giraffe and okapi. *Applied Animal Behaviour Science*, 73 (3), 235-247.
- Bassett, T. and Koli Bi, Z., 1999. Fulbe livestock raising and environmental change in northern Côte d'Ivoire. In: Azarya, V., Breedveld, A., De Bruijn, M., et al. eds. *Pastoralists under pressure? Fulbe societies confronting change in West Africa*. Brill Academic Publishers, Leiden, 139-159.
- Baumont, R., Prache, S., Meuret, M., et al., 2000. How forage characteristics influence behaviour and intake in small ruminants: a review. *Livestock Production Science*, 64 (1), 15-28.
- Bayer, W. and Waters-Bayer, A., 1994. Forage alternatives from range and field: pastoral forage management and improvement in the African drylands. In: Scoones, I. ed. *Living with uncertainty: new directions in pastoral development in Africa*. Intermediate Technology Publications, London, 58-78.
- Beekman, J.H. and Prins, H.H.T., 1989. Feeding strategies of sedentary large herbivores in East Africa with emphasis on the African buffalo, *Syncerus caffer*. *African Journal of Ecology*, 27, 129-147.
- Behnke, R.H. and Scoones, I.E., 1993. Rethinking range ecology: implications for rangeland management in Africa. In: Behnke, R.H. and Scoones, I.E. eds. *Range ecology at disequilibrium: new models of natural variability and pastoral adaptation in Africa savannas*. Overseas Development Institute, London, 1-30.

- Behnke, R.H., Scoones, I.E. and Kerven, C., 1993. *Range ecology at disequilibrium: new models of natural variability and pastoral adaptation in African savannas*. International Livestock Center for Africa ILCA, Addis Ababa.
- Bekure, S., De Leeuw, P.N., Grandin, B.E., et al. (eds.), 1991. *Maasai herding: an analysis of the livestock production system of Maasai pastoralists in eastern Kajiado District, Kenya*. International Livestock Center for Africa ILCA, Addis Ababa. ILCA Systems Study no. 4.
- Bell, R.H.V., 1970. The use of herbaceous layers by grazing ungulates in the Serengeti. In: Watson, A. ed. *Animal populations in relation to their food resources: a symposium of the British Ecological Society, Aberdeen, 24-28 March 1969*. Oxford, Blackwell, 111-124.
- Belovsky, G.E., 1978. Diet optimization in a generalist herbivore: the moose. *Theoretical Population Biology*, 14 (1), 105-134.
- Belovsky, G.E., 1984. Herbivore optimal foraging: a comparative test of three models. *American Naturalist*, 124 (1), 97-115.
- Belovsky, G.E., 1986. Optimal foraging and community structure: implications for a guild of generalist grassland herbivores. *Oecologia*, 70 (1), 35-52.
- Belovsky, G.E. and Schmitz, O.J., 1994. Plant defenses and optimal foraging by mammalian herbivores. *Journal of Mammalogy*, 75 (4), 816-832.
- Belovsky, G.E. and Slade, J.B., 1986. Time budgets of grassland herbivores: body size similarities. *Oecologia*, 70 (1), 53-62.
- Bergman, C.M., Fryxell, J.M. and Gates, C.C., 2000. The effect of tissue complexity and sward height on the functional response of wood bison. *Functional Ecology*, 14 (1), 61-69.
- Bergman, C.M., Fryxell, J.M., Gates, C.C., et al., 2001. Ungulate foraging strategies: energy maximizing or time minimizing? *Journal of Animal Ecology*, 70 (2), 289-300.
- Bernus, E., 1979. Le contrôle du milieu naturel et du troupeau par les éleveurs touaregs sahéliens. In: L'Equipe écologie et anthropologie des sociétés pastorales ed. *Pastoral production and society=Production pastorale et société: proceedings of the international meeting on nomadic pastoralism, Paris 1-3 Dec. 1976*. Cambridge University Press, Cambridge, 67-74.
- Berteaux, D., Crete, M., Huot, J., et al., 1998. Food choice by white-tailed deer in relation to protein and energy content of the diet: a field experiment. *Oecologia*, 115 (1/2), 84-92.
- Biddanda, B., Ogdahl, M. and Cotner, J., 2001. Dominance of bacterial metabolism in oligotrophic relative to eutrophic waters. *Limnology and Oceanography*, 46 (3), 730-739.
- Biondini, M.E., Steuter, A.A. and Hamilton, R.G., 1999. Bison use of fire-managed remnant prairies. *Journal of Range Management*, 52 (5), 454-461.
- Black, J.L. and Kenney, P.A., 1984. Factors affecting diet selection by sheep. 2. Height and density of pasture. *Australian Journal of Agricultural Research*, 35 (4), 565-578.
- Blackman, F.F., 1905. Optima and limiting factors. *Annals of Botany*, 19, 281-295.
- Boice, R., 1972. Some behavioural tests of domestication in Norway rats. *Behaviour*, 42 (3/4), 198-231.
- Bokdam, J. and Wallis De Vries, M.F., 1992. Forage quality as a limiting factor for cattle grazing in isolated Dutch nature reserves. *Conservation Biology*, 6 (3), 399-408.
- Boomker, E.A., 1987. *Fermentation and digestion in the kudu*. PhD Thesis, University of Pretoria, Pretoria.
- Boone, R.B., 2000. *Integrated management and assessment system balancing food security, conservation, and ecosystem integrity in East Africa: training manual*. Global Livestock Collaborative Research Support Program, University of California, Davis. [<http://www.nrel.colostate.edu/projects/imas/prods/rboone/Workshop.html>]
- Boone, R.B., BurnSilver, S.B., Thornton, P.K., et al., 2005. Quantifying declines in livestock due to land subdivision. *Rangeland Ecology and Management*, 58 (5), 523-532.
- Boone, R.B., Coughenour, M.B., Galvin, K.A., et al., 2002. Addressing management questions for Ngorongoro Conservation Area, Tanzania, using the Savanna modelling system. *African Journal of Ecology*, 40 (2), 138-150.
- Boone, R.B., Galvin, K.A., Smith, N.M., et al., 2000. Generalizing El Niño effects upon Maasai livestock using hierarchical clusters of vegetation patterns. *Photogrammetric Engineering and Remote Sensing*, 66 (6), 737-744.
- Boone, R.B. and Hobbs, N.T., 2004. Lines around fragments: effects of fencing on large herbivores. *African Journal of Range and Forage Science*, 21 (3), 147-158.
- Borchers, D.L., Buckland, S.T. and Zucchini, W., 2002. *Estimating animal abundance: closed populations*. Springer, London.
- Bos, D., 2002. *Grazing in coastal grasslands: Brent Geese and facilitation by herbivory*. PhD Thesis, Rijksuniversiteit Groningen, Groningen.
- Boughey, A.S., 1973. *Ecology of populations*. 2nd edn. MacMillan, New York.

- Bowers, M.A. and Brown, J.H., 1982. Structure in a desert rodent community: use of space around *Dipodomys spectabilis* mounds. *Oecologia*, 92 (2), 242-249.
- Box, E.O., Holben, E.N. and Kalb, V., 1989. Accuracy of the AVHRR vegetation index as a predictor of biomass, primary productivity and net CO₂ flux. *Vegetatio*, 80 (2), 71-89.
- Bradbury, J.W., Vehrencamp, S.L., Clifton, K.E., et al., 1996. The relationship between bite rate and local forage abundance in wild Thomson's gazelles. *Ecology*, 77 (7), 2237-2255.
- Braithwaite, L.W., Dudzinski, M.L. and Turner, J., 1983. Studies on the arboreal marsupial fauna of eucalypt forests being harvested for wood pulp at Eden, N.S.W. II. Relationship between the fauna density, richness and diversity, and measured variables of the habitat. *Australian Wildlife Research*, 10 (2), 231-247.
- Breman, H. and De Ridder, N., 1991. *Manuel sur les pâturages des pays sahéliens*. CTA, Wageningen.
- Briske, D.D., Fuhlendorf, S.D. and Smeins, F.E., 2003. Vegetation dynamics on rangelands: a critique of the current paradigms. *Journal of Applied Ecology*, 40 (4), 601-614.
- Brock, B.L. and Owensby, C.E., 2000. Predictive models for grazing distribution: a GIS approach. *Journal of Range Management*, 53 (1), 39-46.
- Brown, J.H., Fox, B.J. and Kelt, D.A., 2000. Assembly rules: desert rodent communities are structured at scales from local to continental. *American Naturalist*, 156 (3), 314-321.
- Brown, J.S., 1999. Vigilance, patch use and habitat selection: foraging under predation risk. *Evolutionary Ecology Research*, 1 (1), 49-71.
- Brown, J.S., Kotler, B.P. and Mitchell, W.A., 1994. Foraging theory, patch use, and the structure of a Negev Desert granivore community. *Ecology*, 75 (8), 2286-2300.
- Brown, J.S., Kotler, B.P., Smith, R.J., et al., 1988. The effects of owl predation on the foraging behavior of heteromyid rodents. *Oecologia*, 76 (3), 408-415.
- Bruce, A. and Hong-Ye, G., 1996. *Applied wavelet analysis with S-plus*. Springer, New York.
- Bryant, J.P. and Kuropat, P.J., 1980. Selection of winter forage by subarctic browsing vertebrates: the role of plant chemistry. *Annual Review of Ecology and Systematics*, 11, 261-285.
- Bryant, J.P., Provenza, F.D., Pastor, J., et al., 1991. Interactions between woody plants and browsing mammals mediated by secondary metabolites. *Annual Review of Ecology and Systematics*, 22, 431-446.
- Buckland, S.T. and Elston, D.A., 1993. Empirical models for the spatial distribution of wildlife. *Journal of Applied Ecology*, 30 (3), 478-495.
- Bunnell, F.L. and Gillingham, M.P., 1985. Foraging behaviour: dynamics of dining out. In: Hudson, R.J. and White, R.G. eds. *Bioenergetics of wild herbivores*. CRC Press, Boca Raton, 53-79.
- Burditt, L., Buchanan, D. and Fitch, J., 1995. *Breeds of livestock: Cattle [website]*. Oklahoma State University, Stillwater. [<http://www.ansi.okstate.edu/breeds/cattle>]
- Burnham, K.P. and Anderson, D.R., 2002. *Model selection and multimodel inference: a practical information-theoretic approach*. Springer, New York.
- BurnSilver, S., Boone, R.B. and Galvin, K.A., 2004. Linking pastoralists to a heterogeneous landscape: the case of four Maasai group ranches in Kajiado District, Kenya. In: Fox, J., Mishra, V., Rindfuss, R., et al. eds. *Linking household and remotely sensed data: methodological and practical problems*. Kluwer Academic Publishing, Boston, 173-199.
- Burritt, E.A. and Provenza, F.D., 2000. Role of toxins in intake of varied diets by sheep. *Journal of Chemical Ecology*, 26 (8), 1991-2005.
- Buttolph, L.P. and Coppock, D.L., 2001. Project Alpaca. *Rangelands*, 23 (2), 10-13.
- Campling, R.C., 1970. Physical regulation of voluntary intake. In: Phillipson, A.J. ed. *Physiology of digestion and metabolism in the ruminant*. Oriel Press, Newcastle, 226-234.
- Caughley, G., 1976. Plant-herbivore systems. In: May, R.M. ed. *Theoretical ecology*. Blackwell, Oxford, 94-113.
- Caughley, G., 1977. Sampling in aerial survey. *Journal of Wildlife Management*, 41 (4), 605-615.
- Chan-McLeod, A.C.A., White, R.G. and Holleman, D.F., 1994. Effects of protein and energy intake, body condition, and season on nutrient partitioning and milk production in caribou and reindeer. *Canadian Journal of Zoology*, 72 (5), 938-947.
- Chang, C., 1993. Pastoral transhumance in the southern Balkans as a social ideology: ethnoarchaeological research in northern Greece. *American Anthropologist*, 95 (3), 687-703.
- Charnov, E.L., 1976. Optimal foraging, the marginal value theorem. *Theoretical Population Biology*, 9 (2), 129-136.
- Chase, J.M. and Leibold, M.A., 2003. *Ecological niches: linking classical and contemporary approaches*. University of Chicago Press, Chicago.

- Chatty, D. and Colchester, M. (eds.), 2002. *Conservation and mobile indigenous peoples: displacement, forced settlement and sustainable development*. Berghahn, New York. Studies in Forced Migration, no. 10.
- Chesson, P., 2000a. General theory of competitive coexistence in spatially-varying environments. *Theoretical Population Biology*, 58 (3), 211-237.
- Chesson, P., 2000b. Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics*, 31, 343-366.
- Cibils, A.F., Howery, L.D. and Ruyle, G.B., 2004. Diet and habitat selection by cattle: the relationship between skin- and gut-defense systems. *Applied Animal Behaviour Science*, 88 (3/4), 187-208.
- Clanet, J.C., 1996. L'évolution récente des mouvements pastorales dans le Bassin Conventionnel du Lac Tchad. In: *CIRAD/CTA Livestock atlas of the Lake Chad basin*. CTA, Wageningen, 97-103.
- Clark, C.W. and Mangel, M., 2000. *Dynamic state variable models in ecology: methods and applications*. Oxford University Press, New York.
- Clark, D.A. and Harris, P.S., 1985. Composition of the diet of sheep grazing swards of differing white clover content and spatial distribution. *New Zealand Journal of Agricultural Research*, 28, 233-240.
- Clarke, J.L., Welch, D. and Gordon, I.J., 1995. The influence of vegetation pattern on the grazing of heather moorland by red deer and sheep. I. The location of animals on grass/heather mosaics. *Journal of Applied Ecology*, 32 (1), 166-176.
- Clauss, M., Frey, R., Kiefer, B., et al., 2003a. The maximum attainable body size of herbivorous mammals: morphophysiological constraints on foregut, and adaptations of hindgut fermenters. *Oecologia*, 136 (1), 14-27.
- Clauss, M., Lason, K., Gehrke, J., et al., 2003b. Captive roe deer (*Capreolus capreolus*) select for low amounts of tannic acid but not quebracho: fluctuation of preferences and potential benefits. *Comparative Biochemistry and Physiology. B. Biochemistry and Molecular Biology*, 136 (2), 369-382.
- Clevers, J.G.P.W., 1988. The derivation of a simplified reflectance model for the estimation of leaf area index. *Remote Sensing of Environment*, 25 (1), 53-69.
- Clevers, J.G.P.W., 1989. The application of a weighted infrared-red vegetation index for estimating leaf area index by correcting for soil moisture. *Remote Sensing of Environment*, 29 (1), 25-37.
- Clevers, J.G.P.W. and Jongschaap, R., 2001. Imaging spectrometry for agricultural applications. In: Van der Meer, F.D. and De Jong, S.M. eds. *Imaging spectrometry: basic principles and prospective applications*. Kluwer Academic Publishers, Dordrecht, 157-199. Remote Sensing and Digital Image Processing no. 4.
- Clevers, J.P.G.W., 1994. Imaging spectroscopy in agriculture: plant vitality and yield indicators. In: Hill, J. and Megier, J. eds. *Imaging spectrometry: a tool for environmental observations*. Kluwer Academic Publishers, Dordrecht, 193-219.
- Coe, M.J., 1983. Large herbivores and food quality. In: Lee, J.A., McNeill, J. and Rorison, I.H. eds. *Nitrogen as an ecological factor: the 22nd symposium of the British Ecological Society, Oxford, 1981*. Blackwell, Oxford, 345-368.
- Coe, M.J., Cumming, D.H. and Phillipson, J., 1976. Biomass and production of large African herbivores in relation to rainfall and primary production. *Oecologia*, 22 (4), 341-354.
- Connell, J.H., 1983. On the prevalence and relative importance of interspecific competition: evidence from field experiments. *American Naturalist*, 122 (5), 661-696.
- Cooper, S.D.B., Kyriazakis, I., Anderson, D.H., et al., 1993. The effect of physiological state (late pregnancy) on the diet selection of ewes. *Animal Production*, 56, 469A.
- Cooper, S.D.B., Kyriazakis, I., Nolan, J.V., et al., 1995. Diet selection in sheep: the role of the rumen environment in the selection of a diet from two feeds that differ in their energy density. *British Journal of Nutrition*, 74 (1), 39-54.
- Cooper, S.M. and Owen-Smith, N., 1986. Effects of plant spinescence on large mammalian herbivores. *Oecologia*, 68 (3), 446-455.
- Cooper, S.M., Owen-Smith, N. and Bryant, J.P., 1988. Foliage acceptability to browsing ruminants in relation to seasonal changes in the leaf chemistry of woody plants in a South African savanna. *Oecologia*, 75 (3), 336-342.
- Coppedge, B.R. and Shaw, J.H., 1998. Bison grazing patterns on seasonally burned tallgrass prairie. *Journal of Range Management*, 51 (3), 258-264.
- Coppock, D.L., Ellis, J.E., Detling, J.K., et al., 1983. Plant-herbivore interactions in a North American mixed-grass prairie. II. Responses of bison to modification of vegetation by prairie dogs. *Oecologia*, 56 (1), 10-15.

- Corsi, F., De Leeuw, J. and Skidmore, A.K., 2000. Modelling species distribution with GIS. *In: Boitani, L. and Fuller, T.K. eds. Research techniques in animal ecology controversies and consequences.* Columbia University Press, New York, 389-434.
- Cosner, C., Deangelis, D.L., Ault, J.S., et al., 1999. Effects of spatial grouping on the functional response of predators. *Theoretical Population Biology*, 56 (1), 65-75.
- Coughenour, M.B., 1991. Spatial components of plant-herbivore interactions in pastoral, ranching, and native ungulate ecosystems. *Journal of Range Management*, 44 (6), 530-542.
- Cowling, R.M., 2000. Challenges to the 'new' rangeland science. *Trends in Ecology and Evolution*, 15 (8), 303-304.
- Crawley, M.J., 1983. *Herbivory : the dynamics of animal-plant interactions.* Blackwell, Oxford. Studies in Ecology no. 10.
- Creel, S., 2001. Four factors modifying the effect of competition on Carnivore population dynamics as illustrated by African wild dogs. *Conservation Biology*, 15 (1), 271-274.
- Critchley, H.D. and Rolls, E.T., 1996. Hunger and satiety modify the responses of olfactory and visual neurons in the primate orbitofrontal cortex. *Journal of Neurophysiology*, 75 (4), 1673-1686.
- Curran, P.J., 1988. The semivariogram in remote sensing: an introduction. *Remote Sensing of Environment*, 24 (3), 493-507.
- Dahl, G. and Hjort, A., 1976. *Having herds: pastoral herd growth and household economy.* University of Stockholm, Stockholm. Stockholm Studies in Social Anthropology no. 2.
- Dale, M.R.T., 1999. *Spatial pattern analysis in plant ecology.* Cambridge University Press, Cambridge.
- Davies, R.A. and Skinner, J.D., 1986. Temporal activity patterns of springbok and merino sheep during a karoo drought. *Transactions of the Royal Society of South Africa*, 46, 133-147.
- Dayan, T. and Simberloff, D., 1994. Character displacement, sexual dimorphism, and morphological variation among British and Irish mustelids. *Ecology*, 75 (4), 1063-1073.
- De Bie, S., 1991. *Wildlife resources of the West African savanna.* PhD Thesis, Wageningen University, Wageningen.
- De Boer, W.F. and Prins, H.H.T., 1990. Large herbivores that strive mightily but eat and drink as friends. *Oecologia*, 82 (2), 264-274.
- De Haan, C., Steinfeld, H. and Blackburn, H., 1997. *Livestock and the environment: finding a balance.* Commission of the European Communities, Brussels.
- De Leeuw, J., Prins, H.H.T., Njuguna, E.C., et al., 1998. *Interpretation of DRSRS: animal counts (1977-1997) in the rangeland districts of Kenya.* ITC, Enschede.
- DeAngelis, D.L., 1992. *Dynamics of nutrient cycling and food webs.* Chapman and Hall, London. Population and Community Biology Series no. 9.
- Delany, M. and Happold, D.C.D., 1979. *Ecology of African mammals.* Longman, London.
- Demment, M.W. and Van Soest, P.J., 1985. A nutritional explanation for body-size patterns of ruminant and nonruminant herbivores. *American Naturalist*, 125 (5), 641-672.
- Denny, P., 1993. Wetlands of Africa: introduction. *In: Whigham, D., Dykyjova, D. and Hejny, S. eds. Wetlands of the world: inventory, ecology and management.* Kluwer, Dordrecht, 1-31. Handbook of Vegetation Science no. 15/2.
- Desforges, M.F. and Wood Gush, D.G.M., 1976. A behavioural comparison of domestic and mallard ducks: habituation and flight reactions. *Animal Behaviour*, 23, 692-697.
- Diamond, J.M., 1975. Assembly of species communities. *In: Cody, M.L. and Diamond, J.M. eds. Ecology and evolution of communities.* Harvard University Press, Cambridge, 342-444.
- Diggle, P.J., 2003. *Statistical analysis of spatial point patterns.* 2nd edn. Academic Press, London.
- Diggle, P.J. and Chetwynd, A.G., 1991. Second-order analysis of spatial clustering for inhomogeneous populations. *Biometrics*, 47 (3), 1155-1163.
- Distel, R.A., Griggs, T.C., Laca, E.A., et al., 1991. Cattle prefer patches that yield greater instantaneous intake rate in patchy ryegrass swards. *In: Proceedings of the second grazing livestock nutrition conference.* Oklahoma State University, 171. MP / Oklahoma State University, Agricultural Experiment Station, Division of Agriculture no. MP-133.
- Distel, R.A., Laca, E.A., Griggs, T.C., et al., 1995. Patch selection by cattle: maximization of intake rate in horizontally heterogeneous pastures. *Applied Animal Behaviour Science*, 45 (1/2), 11-21.
- Drake, J.A., Zimmermann, C.R., Purucker, T., et al., 1999. On the nature of the assembly trajectory. *In: Weiher, E. and Keddy, P. eds. Ecological assembly rules: perspectives, advances, retreats.* Cambridge University Press, Cambridge, 233-250.
- Drent, R.H. and Prins, H.H.T., 1987. The herbivore as prisoner of its food supply. *In: Van Andel, J., Bakker, J. and Snaydon, R.W. eds. Disturbance in grasslands: species and population responses.* Dr. W. Junk Publishing Company, Dordrecht, 133-149.

- Drescher, M., 2003. *Grasping complex matter: large herbivore foraging in patches of heterogeneous resources*. PhD Thesis, Wageningen University.
- Drescher, M., Heitkönig, I.M.A., Raats, J.G., et al., 2006a. The role of grass stems as structural foraging deterrents and their effects on the foraging behaviour of cattle. *Applied Animal Behaviour Science*, 101 (1/2), 10-26.
- Drescher, M., Heitkönig, I.M.A., Van den Brink, P.J., et al., 2006b. Effects of sward structure on herbivore foraging behaviour in a South African savanna: an investigation of the forage maturation hypothesis. *Austral Ecology*, 31 (1), 76-87.
- Du Toit, J.G., 1996. Water requirements. In: Bothma, J.d.P. ed. *Game ranch management*. 3rd edn. Van Schaik JL, Pretoria, 95-98.
- Dublin, H., 1995. Vegetation dynamics in the Serengeti-Mara ecosystem: the role of elephants, fire, and other factors. In: Sinclair, A.R.E. and Arcese, P. eds. *Serengeti 2: dynamics, management, and conservation of an ecosystem [based on papers at a workshop held in December 1991 at the Serengeti Research Institute, Tanzania]*. University of Chicago Press, Chicago, 71-90.
- Dublin, H.T., Sinclair, A.R.E. and McGlade, J., 1990. Elephants and fire as causes of multiple stable states in the Serengeti-Mara woodlands. *Journal of Animal Ecology*, 59 (3), 1147-1164.
- Dumont, B. and Boissy, A., 2000. Grazing behaviour of sheep in a situation of conflict between feeding and social motivations. *Behavioural Processes*, 49 (3), 131-138.
- Dumont, B., Maillard, J.F. and Petit, M., 2000. The effect of the spatial distribution of plant species within the sward on the searching success of sheep when grazing. *Grass and Forage Science*, 55 (2), 138-145.
- Dumont, B. and Petit, M., 1998. Spatial memory of sheep at pasture. *Applied Animal Behaviour Science*, 60 (1), 43-53.
- Dunbrack, R.L. and Ramsay, A., 1993. The allometry of mammalian adaptations to seasonal environments: a critique of the fasting endurance hypothesis. *Oikos*, 66 (2), 336-342.
- Duncan, A.J. and Gordon, I.J., 1999. Habitat selection according to the ability of animals to eat, digest and detoxify foods. *Proceedings of the Nutrition Society*, 58 (4), 799-805.
- Duncan, A.J., Hartley, S.E. and Iason, G.R., 1998. The effect of previous browsing damage on the morphology and chemical composition of Sitka spruce (*Picea sitchensis*) saplings and on their subsequent susceptibility to browsing by red deer (*Cervus elaphus*). *Forest Ecology and Management*, 103 (1), 57-67.
- Duncan, A.J., Mayes, R.W., Young, S.A., et al., 2001. Choice of foraging patches by hill sheep given different opportunities to seek shelter and food. *Animal Science*, 73 (3), 563-570.
- Duncan, A.J., Reid, S.A., Thoss, V., et al., 2005. Browse selection in response to simulated seasonal changes in diet quality through postingestive effects. *Journal of Chemical Ecology*, 31 (4), 729-744.
- Duncan, P., 1975. *Topi and their food supply*. PhD Thesis, University of Nairobi, Nairobi.
- Duncan, P., 1983. Determinants of the use of habitat by horses in a Mediterranean wetland. *Journal of Animal Ecology*, 52, 93-109.
- Duncan, P., 1985. Time budgets of Camargue horses. III. Environmental influences. *Behaviour*, 92, 188-208.
- Durant, S.M., 1998. Competition refuges and coexistence: an example from Serengeti carnivores. *Journal of Animal Ecology*, 67 (3), 370-386.
- Dury, S., Turner, B., Foley, B., et al., 2001. Use of high spectral resolution remote sensing to determine leaf palatability of eucalypt trees for folivorous marsupials. *International Journal of Applied Earth Observation and Geoinformation*, 3 (4), 328-336.
- Dyson-Hudson, N., 1980. Strategies of resource exploitation among East African savanna pastoralists. In: Harris, D.R. ed. *Human ecology in savanna environments*. Academic Press, London, 171-184.
- Dyson-Hudson, R. and Dyson-Hudson, N., 1980. Nomadic pastoralism (Africa). *Annual review of anthropology*, 9, 15-61.
- Early, D.M. and Provenza, F.D., 1998. Food flavour and nutritional characteristics alter dynamics of food preference in lambs. *Journal of Animal Science*, 76 (3), 728-734.
- East, R., 1984. Rainfall, soil nutrient status and biomass of large African savanna mammals. *African Journal of Ecology*, 22 (4), 245-270.
- Edwards, G.R., Newman, J.A., Parsons, A.J., et al., 1996. The use of spatial memory by grazing animals to locate food patches in spatially heterogeneous environments: an example with sheep. *Applied Animal Behaviour Science*, 50 (2), 147-160.
- Egan, A.R., 1980. Host animal-rumen relationships. *Proceedings of the Nutrition Society*, 39 (1), 79-87.
- Eklundh, L., 1998. Estimating relations between AVHRR NDVI and rainfall in East Africa at 10-day and monthly time scales. *International Journal of Remote Sensing*, 19 (3), 563-568.

- Ellenbroek, G.A., 1987. *Ecology and productivity of an African wetland system: the Kafue Flats, Zambia*. Junk, Dordrecht. Geobotany no. 9.
- Ellis, J.E., 1995. Climate variability and complex ecosystem dynamics: implications for pastoral development. In: Scoones, I. ed. *Living with uncertainty: new directions in pastoral development in Africa*. Intermediate Technology Publications, London, 37-46.
- Ellis, J.E. and Coughenour, M.B., 1998. The SAVANNA integrated modelling system: an integrated remote sensing, GIS and spatial simulation modelling approach. In: Squires, V.R. and Sidahmed, A.E. eds. *Drylands: sustainable use of rangelands into the twenty-first century*. IFAD, Rome, 97-106.
- Ellis, J.E. and Swift, D.M., 1988. Stability of African pastoral ecosystems: alternate paradigms and implications for development. *Journal of Range Management*, 41 (6), 450-459.
- Elton, C., 1927. *Animal ecology*. Sidgwick and Jackson, London.
- Elvidge, C.D., 1990. Visible and near infrared reflectance characteristics of dry plant materials. *International Journal of Remote Sensing*, 11 (10), 1775-1795.
- Elvidge, C.D., Zhikang, C. and Groeneveld, D.P., 1993. Detection of trace quantities of green vegetation in 1990 AVIRIS data. *Remote Sensing of Environment*, 44 (2/3), 271-279.
- Engel, C., 2002. *Wild health: how animals keep themselves well and what we can learn from them*. Houghton Mifflin Co., New York.
- Enkh-Amgalan, A., 2002. Change and sustainability of pastoral land use systems in Mongolia. In: Chuluun, T. and Ojima, D. eds. *Fundamental issues affecting sustainability of the Mongolian steppe*. International Institute for Study of Nomadic Civilizations, Ulaanbaatar, 228-231.
- Enkhtuvshin, B. and Tumujav, M., 2002. World nomads and Mongolian traditional pasture use. In: Chuluun, T. and Ojima, D. eds. *Fundamental issues affecting sustainability of the Mongolian steppe*. International Institute for Study of Nomadic Civilizations, Ulaanbaatar, 19-25.
- Estes, J.E. and Mooneyhan, D.W., 1994. Of maps and myths. *Photogrammetric Engineering and Remote Sensing*, 60 (5), 517-524.
- Etzenhouser, M.J., Owens, M.K., Spalinger, D.E., et al., 1998. Foraging behaviour of browsing ruminants in a heterogeneous landscape. *Landscape Ecology*, 13 (1), 55-64.
- Evans-Pritchard, E.E., 1940. *The nuer: a description of the modes of livelihood and political institutions of a Nilotic people*. Oxford University Press, Oxford.
- Evans, M., Green, B. and Newgrain, K., 2003. The field energetics and water fluxes of free-living wombats (Marsupialia: Vombatidae). *Oecologia*, 137 (2), 171-180.
- Fancy, S.G. and White, R.G., 1985. Incremental cost of activity. In: Hudson, R.J. and White, R.G. eds. *Bioenergetics of wild herbivores*. CRC Press, Boca Raton, 143-159.
- FAO, 2001. *Pastoralism in the new millenium*. FAO, Rome. FAO Animal Production and Health Paper no. 150.
- Farnsworth, K.D. and Beecham, J.A., 1999. How do grazers achieve their distribution? A continuum of models from random diffusion to the ideal free distribution using biased random walks. *American Naturalist*, 153 (5), 509-526.
- Farnsworth, K.D. and Illius, A.W., 1996. Large grazers back in the fold: generalizing the prey model to incorporate mammalian herbivores. *Functional Ecology*, 10 (5), 678-680.
- Ferguson, S.H., Bergerud, A.T. and Ferguson, R., 1988. Predation risk and habitat selection in the persistence of a remnant caribou population. *Oecologia*, 76 (2), 236-245.
- Fernandez-Gimenez, M.E. and Allen-Diaz, B., 1999. Testing a non-equilibrium model of rangeland vegetation dynamics in Mongolia. *Journal of Applied Ecology*, 36 (6), 871-885.
- Ferwerda, J.G., 2005. *Charting the quality of forage: measuring and mapping the variation of chemical components in foliage with hyperspectral remote sensing*. PhD Thesis, Wageningen University, Wageningen.
- Ferwerda, J.G., Skidmore, A.K., Van Wieren, S.E., et al., 2002. Field spectrometry of leaves for nutrient assesment. In: Navalgund, R.R., Nayak, S.R., Sudarshana, R., et al. eds. *ISPRS 2002 : proceedings of ISPRS commission VII international symposium : resource and environmental monitoring, December 3-6 2002, Hyderabad, India*. National Remote Sensing Agency, Hyderabad.
- Feyaerts, F. and Van Gool, L., 2001. Multi-spectral vision system for weed detection. *Pattern Recognition Letters*, 22 (6/7), 667-674.
- Field, C. and Mooney, H.A., 1986. The photosynthesis-nitrogen relationship in wild plants. In: Givnish, T.J. ed. *On the economy of plant form and function: proceedings of the sixth Maria Moors Cabot symposium, "Evolutionary constraints on primary productivity: adaptive patterns of energy capture in plants," Harvard Forest, August 1983*. Cambridge University Press, Cambridge, 25-55.
- Field, C.R., 1976. Palatability factors and nutritive values of the food of buffaloes (*Syncerus caffer*) in Uganda. *East African Wildlife Journal*, 14, 181-201.

- Fisher, D.S., 1996. Modeling ruminant feed intake with protein, chemostatic, and distention feedbacks. *Journal of Animal Science*, 74 (12), 3076-3081.
- Focardi, S. and Marcellini, P., 1995. A mathematical framework for optimal foraging of herbivores. *Journal of Mathematical Biology*, 33 (4), 365-387.
- Focardi, S., Marcellini, P. and Montanaro, P., 1996. Do ungulates exhibit a food density threshold? A field study of optimal foraging and movement patterns. *Journal of Animal Ecology*, 65 (5), 606-620.
- Gordon, I.J., 2003. Browsing and grazing ruminants: are they different beasts? *Forest Ecology and Management*, 181 (1/2), 13-21.
- Gordon, I.J., Duncan, P., Grillas, P., et al., 1990. The use of domestic herbivores in the conservation of the biological richness of European wetlands. *Bulletin d'Ecologie*, 21 (3), 49-60.
- Gordon, I.J. and Illius, A.W., 1988. Incisor arcade structure and diet selection in ruminants. *Functional Ecology*, 2, 15-22.
- Gordon, I.J. and Illius, A.W., 1994. The functional significance of the browser-grazer dichotomy in African ruminants. *Oecologia*, 98 (2), 167-175.
- Gordon, I.J. and Illius, A.W., 1996. The nutritional ecology of African ruminants: a reinterpretation. *Journal of Animal Ecology*, 65 (1), 18-28.
- Gordon, I.J., Illius, A.W. and Milne, J.D., 1996. Sources of variation in the foraging efficiency of grazing ruminants. *Functional Ecology*, 10 (2), 219-226.
- Grace, J.B., 1995. On the measurement of plant competition intensity. *Ecology*, 76 (1), 305-308.
- Graham, R.W. and Lundelius, E.L., 1984. Coevolutionary disequilibrium and Pleistocene extinctions. In: Martin, P.S. and Klein, R.G. eds. *Quaternary extinctions: a prehistoric revolution*. University of Arizona Press, Tucson, 223-249.
- Greenwood, P.T. and Rittenhouse, L.R., 1997. Feeding area selection: the leader-follower phenomena. *Proceedings of the Western Section American Society of Animal Science*, 48, 267-269.
- Grenfell, B.T., Wilson, K., Finkenstadt, B.F., et al., 1998. Noise and determinism in synchronized sheep dynamics. *Nature*, 394 (6694), 674-677.
- Grinnell, J., 1917. The niche-relationships of the California thrasher. *Auk*, 34, 427-433.
- Groombridge, B., 1992. *Global biodiversity: status of the earth's living resources*. Chapman and Hall, London.
- Groot-Bruinderink, G.W.T.A., Lammertsma, D.R., Kramer, K., et al., 1999. *Dynamische interacties tussen hoefdieren en vegetatie in de Oostvaardersplassen*. IBN-DLO, Wageningen. IBN Rapport no. 436.
- Gross, J.E., Hobbs, N.T. and Wunder, B.A., 1993a. Independent variables for predicting intake rate of mammalian herbivores: biomass density, plant density, or bite size? *Oikos*, 68 (1), 75-81.
- Gross, J.E., Shipley, L.A., Hobbs, N.T., et al., 1993b. Functional response of herbivores in food-concentrated patches: tests of a mechanistic model. *Ecology*, 74 (3), 778-791.
- Gross, J.E., Wang, Z. and Wunder, B.A., 1985. Effects of food quality and energy needs: changes in gut morphology and capacity of *Microtus ochrogaster*. *Journal of Mammalogy*, 66 (6), 661-667.
- Gross, J.E., Zank, C., Hobbs, N.T., et al., 1995. Movement rules for herbivores in spatially heterogeneous environments: responses to small scale pattern. *Landscape Ecology*, 10 (4), 209-217.
- Grünbaum, D., 1998. Using spatially explicit models to characterize foraging performance in heterogeneous landscapes. *American Naturalist*, 151 (2), 97-115.
- Grunblatt, J., Said, M.Y., Wargute, P., et al., 1995. *DRSRS aerial surveys database*. Department of Resource Surveys and Remote Sensing, Ministry of Planning and National Development, Nairobi.
- Gustafsson, M., Jensen, P., De Jonge, F.H., et al., 1999. Domestication effects on foraging strategies in pigs (*Sus scrofa*). *Applied Animal Behaviour Science*, 62 (4), 305-317.
- Guthrie, R.D., 1990. *Frozen fauna of the mammoth steppe: the story of Blue Babe*. University of Chicago Press, Chicago.
- Guyot, G., Baret, F. and Major, D.J., 1988. High spectral resolution: determination of spectral shifts between the red and near infrared. *International Archives of Photogrammetry and Remote Sensing*, 27, 750-760.
- Hagerman, A.E., Riedl, K.M., Jones, G.A., et al., 1998. High molecular weight plant polyphenolics (tannins) as biological antioxidants. *Journal of Agricultural and Food Chemistry*, 46 (5), 1887-1892.
- Haltenorth, T. and Diller, H., 1994. *Mammals of Africa including Madagascar*. Harper Collins Publishers, London.
- Hanski, I., 1998. Metapopulation dynamics. *Nature*, 396 (6706), 41-49.
- Hanski, I., 1999. *Metapopulation ecology*. Oxford University Press, Oxford.
- Hardin, G., 1968. The tragedy of the commons: the population problem has no technical solution; it requires a fundamental extension in morality. *Science*, 162 (859), 1243-1248.

- Harrington, R., Owen-Smith, N., Viljoen, P.C., et al., 1999. Establishing the causes of the roan antelope decline in the Kruger National Park, South Africa. *Biological Conservation*, 90 (1), 69-78.
- Harris, N.R., Johnson, D.E., George, M.R., et al., 2002. The effect of topography, vegetation, and weather on cattle distribution at the San Joaquin Experimental Range, California. In: Standiford, R.B., McCreary, D. and Purcell, K.L. eds. *Proceedings of the fifth symposium on oak woodlands: oaks in California's changing landscape. 2001 October 22-25; San Diego, CA*. USDA Forest Service, Albany, 53-63. General Technical Report no. PSW-GTR-184.
- Hary, I., Schwartz, H.J., Pielert, V.H.C., et al., 1996. Land degradation in African pastoral systems and the destocking controversy. *Ecological Modelling*, 86 (2/3), 227-233.
- Haskell, J.P., Ritchie, M.E. and Olff, H., 2002. Fractal geometry predicts varying body size scaling relationships for mammal and bird home ranges. *Nature*, 418 (6897), 527-530.
- Haxeltine, A., 1996. *Modelling the vegetation of the earth*. PhD Thesis, Lund University, Department of Ecology, Lund.
- Heasley, L. and Delehanty, J., 1996. The politics of manure: resource tenure and the agropastoral economy in Southwestern Niger. *Society and Natural Resources*, 9 (1), 31-46.
- Heath, B., 2000. Ranching: an economic yardstick. In: Prins, H.H.T. and Grootenhuys, J.G. eds. *Wildlife conservation by sustainable use*. Kluwer Academic Publishers, Boston, 21-33.
- Hebblewhite, M. and Pletscher, D.H., 2002. Effects of elk group size on predation by wolves. *Canadian Journal of Zoology*, 80 (5), 800-809.
- Hebblewhite, M., Pletscher, D.H. and Paquet, P.C., 2002. Elk population dynamics in areas with and without predation by recolonizing wolves in Banff National Park, Alberta. *Canadian Journal of Zoology*, 80 (5), 789-799.
- Herbel, C.H. and Nelson, A.B., 1966. Activities of Hereford and Santa Gertrudis cattle on a southern New Mexico range. *Journal of Range Management*, 19 (4), 173-176.
- Hernes, P.J. and Hedges, J.I., 2000. Determination of condensed tannin monomers in environmental samples by capillary gas chromatography of acid depolymerization extracts. *Analytical Chemistry*, 72 (20), 5115-5124.
- Hester, A.J., Gordon, I.J., Baillie, G.J., et al., 1999. Foraging behaviour of sheep and red deer within natural heather/grass mosaics. *Journal of Applied Ecology*, 36 (1), 133-146.
- Hiernaux, P. and Diarra, L., 1983. Pâturages de la zone d'inondation du Niger. In: Wilson, R.T., De Leeuw, P.N. and De Haan, C. eds. *Recherches sur les systèmes des zones arides du Mali: résultats préliminaires*. International Livestock Center for Africa ILCA, Addis Abeba, 42-48. Rapport de Recherche no. 5.
- Hik, D.S., 1995. Does risk of predation influence population dynamics? Evidence from the cyclic decline of snowshoe hares. *Wildlife Research*, 22 (1), 115-129.
- Hirst, S.M., 1969. Populations in a Transvaal lowland nature reserve. *Zoologica Africana*, 4, 199-230.
- Hobbs, N.T., 1990. Diet selection by generalist herbivores: a test of the linear programming model. In: Hughes, R.N. ed. *Behavioural mechanisms of food selection*. Springer, Berlin, 395-413.
- Hobbs, N.T., Gross, J.E., Shipley, L.A., et al., 2003. Herbivore functional response in heterogeneous environments: a contest among models. *Ecology*, 84 (3), 666-681.
- Hodgson, J., Cosgrove, G.P. and Woodward, S.J.R., 1997. Research on foraging strategy: progress and priorities. In: Buchanan-Smith, J.G., Bailey, L.D. and McCaughey, P. eds. *Proceedings of the XVIII international grassland congress, Saskatoon, Saskatchewan, Canada, 8-13 June 1997*. Association Management Centre, Calgary, 109-118.
- Hodgson, J., Niezen, J.H., Montossi, F., et al., 1996. Comparative studies on animal performance and parasite infestation in sheep grazing Yorkshire fog, perennial ryegrass, and tall fescue pastures. *Proceedings of the New Zealand Grassland Association*, 57, 89-93.
- Hofmann, R.R., 1973. *The ruminant stomach: stomach structure and feeding habits of East African game ruminants*. East African Literature Bureau, Nairobi. East African Monographs in Biology no. 2.
- Hofmann, R.R., 1989. Evolutionary steps of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia*, 78 (4), 443-457.
- Hofmann, R.R. and Stewart, D.R.M., 1972. Grazer or browser: a classification based on the stomach structure and feeding habits of East African ruminants. *Mammalia*, 36 (2), 226-240.
- Holdridge, L.R., 1947. Determination of world plant formations from simple climatic data. *Science*, 105 (2727), 367-368.
- Holechek, J.L., Pieper, R.D. and Herbel, C.H., 2001. *Range management: principles and practices*. 4th edn. Prentice Hall, Englewood Cliffs.
- Holling, C.S., 1959. Some characteristics of simple types of predation and parasitism. *Canadian Entomologist*, 91 (7), 385-398.

- Homewood, K.M. and Rodgers, W.A., 1991. *Maasailand ecology: pastoralist development and wildlife conservation in Ngorongoro, Tanzania*. Cambridge Univ. Press, Cambridge.
- Hooper, J.F., Workman, J.P., Grumbles, J.B., et al., 1969. Improved livestock distribution with fertilizer: a preliminary economic evaluation. *Journal of Range Management*, 22, 108-110.
- Houseal, G.A. and Olson, B.E., 1995. Cattle use of microclimates on a northern latitude winter range. *Canadian Journal of Animal Science*, 75, 501-507.
- Houston, A. and McNamara, J.M., 1999. *Models of adaptive behaviour: an approach based on state*. Cambridge University Press, Cambridge.
- Howell, P., Lock, M. and Cobb, S., 1988. *The Jonglei Canal: impact and opportunity*. Cambridge University Press, Cambridge.
- Howery, L.D., Bailey, D.W., Ruyle, G.B., et al., 2000. Cattle use visual cues to track food locations. *Applied Animal Behaviour Science*, 67 (1/2), 1-14.
- Howery, L.D., Provenza, F.D., Banner, R.E., et al., 1996. Differences in home range and habitat use among individuals in a cattle herd. *Applied Animal Behaviour Science*, 49 (3), 305-320.
- Howery, L.D., Provenza, F.D., Banner, R.E., et al., 1998. Social and environmental factors influence cattle distribution on rangeland. *Applied Animal Behaviour Science*, 55 (3/4), 231-244.
- Hubbell, S.P., 2001. *The unified neutral theory of biodiversity and biogeography*. Princeton University Press, Princeton. Monographs in Population Biology no. 32.
- Hudson, R.J., 1985. Body size, energetics, and adaptive radiation. In: Hudson, R.J. and White, R.G. eds. *Bioenergetics of wild herbivores*. CRC Press, Boca Raton, 1-24.
- Hudson, R.J. and Frank, S., 1987. Foraging ecology of bison in aspen boreal habitats. *Journal of Range Management*, 40 (1), 71-75.
- Huggard, D.J., 1994. A linear programming model of herbivore foraging: imprecise, yet successful? *Oecologia*, 100 (4), 470-474.
- Humphrey, C. and Sneath, D., 1999. *The end of nomadism? Society, state and the environment in inner Asia*. White Horse Press, Cambridge.
- Hunter, R.F., 1962. Hill sheep and their pasture: a study of sheep-grazing in south-east Scotland. *The Journal of Ecology*, 50 (3), 651-680.
- Huntingford, G.W.B., 1933. Some aspects of Nandi stock-raising. *Journal of the East Africa and Uganda Natural History Society*, 11, 250-262.
- Huntingford, G.W.B., 1953a. *The northern Nilo-Hamites*. International African Institute, London.
- Huntingford, G.W.B., 1953b. *The southern Nilo-Hamites*. International African Institute, London.
- Hutchings, M.J., 1986. The structure of plant populations. In: Crawley, M.J. ed. *Plant ecology*. Blackwell, Oxford, 97-136.
- Hutchins, M., Hancocks, D. and Crockett, C., 1984. Naturalistic solutions to the behavioral problems of captive animals. *Der Zoologische Garten*, 54, 28-42.
- Hutchinson, G.E., 1957. Concluding remarks: Cold Spring Harbor symposium. *Quantitative Biology*, 22, 415-427.
- Iason, G.R., Mantecon, A.R., Sim, D.A., et al., 1999. Can grazing sheep compensate for a daily foraging time constraint? *Journal of Animal Ecology*, 68 (1), 87-93.
- Iason, G.R. and Van Wieren, S.E., 1999. Digestive and ingestive adaptations of mammalian herbivores to low quality forage. In: Olf, H., Brown, V.K. and Drent, R.H. eds. *Herbivores: between plants and predators: the 38th symposium of the British Ecological Society in cooperation with the Netherlands Ecological Society, held at Wageningen Agricultural University, The Netherlands, 1997*. Blackwell Scientific Publications, Oxford, 337-369.
- ILCA, 1979. *Livestock production in the subhumid zone of West Africa: a regional review*. International Livestock Center for Africa ILCA, Addis Ababa. ILCA Systems Study no. 2.
- Illius, A.W. and Gordon, I.J., 1987. The allometry of food intake in grazing ruminants. *Journal of Animal Ecology*, 56 (3), 989-999.
- Illius, A.W. and Gordon, I.J., 1991. Prediction of intake and digestion in ruminants by a model of rumen kinetics integrating animal size and plant characteristics. *Journal of Agricultural Science*, 116 (1), 145-157.
- Illius, A.W. and Gordon, I.J., 1992. Modelling the nutritional ecology of ungulate herbivores: evolution of body size and competitive interactions. *Oecologia*, 89 (3), 428-434.
- Illius, A.W. and O'Connor, T.G., 1999. On the relevance of nonequilibrium concepts to arid and semiarid grazing systems. *Ecological Applications*, 9 (3), 798-813.
- Illius, A.W. and O'Connor, T.G., 2000. Resource heterogeneity and ungulate population dynamics. *Oikos*, 89 (2), 283-294.

- Janis, C.M., 1995. Correlation between craniodental morphology and feeding behavior in ungulates: reciprocal illumination between living and fossil taxa. *In: Thomason, J.J. ed. Functional morphology in vertebrate paleontology*. Cambridge University Press, Cambridge, 76-98.
- Janis, C.M. and Ehrhardt, D., 1988. Correlation of relative muzzle width and relative incisor width with dietary preference in ungulates. *Zoological Journal of the Linnean Society*, 92 (3), 267-284.
- Jarman, M.V. and Jarman, P.J., 1973. Daily activity of impala. *East African Wildlife Journal*, 11, 75-92.
- Jarman, P.J., 1974. The social organization of antelope in relation to their ecology. *Behaviour*, 48 (2), 15-66.
- Jarman, P.J. and Sinclair, A.R.E., 1979. Feeding strategy and pattern of resource partitioning in ungulates. *In: Sinclair, A.R.E. and Norton-Griffiths, M. eds. Serengeti: dynamics of an ecosystem*. University of Chicago Press, Chicago, 130-160.
- Jensen, S.P. and Doncaster, C.P., 1999. Lethal toxins in non-preferred foods: how plant chemical defences can drive microtine cycles. *Journal of Theoretical Biology*, 199 (1), 63-85.
- Jetz, W., Carbone, C., Fulford, J., et al., 2004. The scaling of animal space use. *Science*, 306 (5694), 266-268.
- Jiang, Z. and Hudson, R.J., 1993. Optimal grazing of wapiti (*Cervus elaphus*) on grassland: patch and feeding station departure rules. *Evolutionary Ecology*, 7 (5), 488-498.
- Jina, P.S., 1999. The Chang Pa of the Ladakh Himalayas: some observations on their economy and environment. *Nomadic Peoples*, 3 (1), 94-105.
- Johnson, B.J., Fogel, M.L. and Miller, G.H., 1998. Stable isotopes in modern ostrich eggshell: a calibration for paleoenvironmental applications in semi-arid regions of southern Africa. *Geochimica et Cosmochimica Acta*, 62 (14), 2451-2461.
- Johnson, C.J., Parker, K.L. and Heard, D.C., 2001. Foraging across a variable landscape: behavioral decisions made by woodland caribou at multiple spatial scales. *Oecologia*, 127 (4), 590-602.
- Johnson, C.J., Parker, K.L., Heard, D.C., et al., 2002a. Movement parameters of ungulates and scale-specific responses to the environment. *Journal of Animal Ecology*, 71 (2), 225-235.
- Johnson, C.J., Parker, K.L., Heard, D.C., et al., 2002b. A multiscale behavioral approach to understanding the movements of woodland caribou. *Ecological Applications*, 12 (6), 1840-1860.
- Jones, R.J., Meyer, J.H.F., Bechaz, F.M., et al., 2001. Comparison of rumen fluid from South African game species and from sheep to digest tanniniferous browse. *Australian Journal of Agricultural Research*, 52 (4), 453-460.
- Kalliola, R. and Syrjanen, K., 1991. To what extent are vegetation types visible in satellite imagery? *Annales Botanici Fennici*, 28 (1), 45-57.
- Kaluzny, S.P., Vega, S.C., Cardoso, T.P., et al., 1998. *S+SpatialStats: user's manual for Windows and UNIX*. Springer, New York.
- Kari, S. and Scholte, P., 2001. La réhabilitation pastorale de la plaine d'inondation Waza-Logone (Cameroun): comment consolider sa réussite écologique? *In: Tielkes, E. and Schlecht, E. eds. Elevage et gestion de parcours au Sahel, implications pour le développement: comptes-rendus d'un atelier régional ouest-africain sur le thème "la gestion des pâturages et les projets de développement: quelles perspectives?", tenu du 2 au 6 octobre 2000 à Niamey, Niger*. Grauer, Beuren, 315-316.
- Kavoori, P.S., 1999. *Pastoralism in expansion: the transhuming herders of western Rajasthan*. Oxford University Press, Oxford.
- Kelt, D.A. and Martínez, D.R., 1989. Notes on distribution and ecology of two marsupials endemic to the Valdivian forests of southern South America. *Journal of Mammalogy*, 70 (1), 220-224.
- Kelt, D.A., Rogovin, K., Shembrot, G., et al., 1999. Patterns in the structure of Asian and North American desert small mammal communities. *Journal of Biogeography*, 26 (4), 825-841.
- Kennedy, M. and Gray, R.D., 1993. Can ecological theory predict the distribution of foraging animals? A critical analysis of experiments on the ideal free distribution. *Oikos*, 68 (1), 158-166.
- Kerven, C., 2002. The loss of livestock mobility in the new market economies of Kazakstan and Turkmenistan: implications for range degradation and regeneration. *In: Chuluun, T. and Ojima, D. eds. Fundamental issues affecting sustainability of the Mongolian steppe*. International Institute for Study of Nomadic Civilizations, Ulaanbaatar, 34-51.
- Kerven, C., Alimaev, I.I., Behnke, R., et al., 2004. Retraction and expansion of flock mobility in Central Asia: costs and consequences. *In: Allsopp, N., Palmer, A.R., Milton, S.J., et al. eds. Proceedings of the VII International Rangelands Congress, Durban South Africa, 26 July-1 August 2003*. 543-556.
- Key, C. and MacIver, R.M., 1980. The effects of maternal influences on sheep: breed differences in grazing, resting and courtship behaviour. *Applied Animal Ethology*, 6, 33-48.

- Khaemba, W.M. and Stein, A., 2000. Use of GIS for a spatial and temporal analysis of Kenyan wildlife with generalised linear modelling. *International Journal of Geographical Information Science*, 14 (8), 833-853.
- Kijazi, A., Mkumbo, S. and Thompson, D.M., 1997. Human and livestock population trends. In: Thompson, D.M. ed. *Multiple land-use: the experience of the Ngorongoro conservation area, Tanzania*. IUCN, Gland, 169-180.
- Kimani, K. and Pickard, J., 1998. Recent trends and implications of group ranch sub-division and fragmentation in Kajiado district, Kenya. *The Geographical Journal*, 164 (2), 202-213.
- Kingdon, J., 1982. *East African mammals: an atlas of evolution in Africa. Vol. 3, pt. C: Bovids*. Academic Press, London.
- Kingdon, J., 1997. *The Kingdon field guide to African mammals*. Academic Press, San Diego.
- Klein, D.R. and Fairall, N., 1986. Comparative foraging behaviour and associated energetics of impala and blesbok. *Journal of Applied Ecology*, 23 (2), 489-502.
- Knipling, E.B., 1970. Physical and physiological basis for the reflectance of visible and near-infrared radiation. *Remote Sensing of Environment*, 1, 155-159.
- Kolasa, J. and Pickett, S.T.A. (eds.), 1991. *Ecological heterogeneity*. Springer, Berlin. Ecological Studies, Analysis and Synthesis no. 86.
- Kondo, M., Kita, K. and Yokota, H.O., 2004. Feeding value to goats of whole-crop oat ensiled with green tea waste. *Animal Feed Science and Technology*, 113 (1/4), 71-81.
- Köppen, W., 1931. *Grundriss der Klimakunde*. Willem de Gruyter, Berlin.
- Kotler, B.P. and Brown, J.S., 1999. Mechanisms of coexistence of optimal foragers as determinants of local abundances and distributions of desert granivores. *Journal of Mammalogy*, 80 (2), 361-374.
- Kotliar, N.B. and Wiens, J.A., 1990. Multiple scales of patchiness and patch structure: a hierarchical framework for the study of heterogeneity. *Oikos*, 59 (2), 253-260.
- Kramer, K., Groot Bruinderink, G.W.T.A. and Prins, H.H.T., 2006. Spatial interactions between ungulate herbivory and forest management. *Forest Ecology and Management*, 226 (1/3), 238-247.
- Kraus, T.E.C., Zasoski, R.J., Dahlgren, R.A., et al., 2004. Carbon and nitrogen dynamics in a forest soil amended with purified tannins from different plant species. *Soil Biology and Biochemistry*, 36 (2), 309-321.
- Krause, J. and Ruxton, G.D., 2002. *Living in Groups*. Oxford University Press, Oxford.
- Krebs, C.J., 1989. *Ecological methodology*. Harper Collins, New York.
- Krebs, J.R. and Davies, N.B. (eds.), 1997. *Behavioural ecology: an evolutionary approach*. 2nd edn. Blackwell, Oxford.
- Krebs, J.R., Ryan, J.C. and Charnov, E.L., 1974. Hunting by expectation or optimal foraging? A study of patch use by chickadees. *Animal Behaviour*, 22 (4), 953-964.
- Kristjanson, P., Radeny, M., Nkedianye, D., et al., 2002. *Valuing alternative land-use options in the Kitengela wildlife dispersal area of Kenya*. International Livestock Research Institute ILRI, Nairobi. ILRI Impact Assessment Series no. 10.
- Křivan, V. and Vrkoč, I., 2000. Patch choice under predation hazard. *Theoretical Population Biology*, 58 (4), 329-340.
- Ksikisi, T. and Laca, E.A., 2000. Can social interactions affect food searching efficiency of cattle? *Rangeland Journal*, 22 (2), 235-242.
- Kumar, L., Schmidt, K.S., Dury, S., et al., 2001. Imaging spectroscopy and vegetation science. In: Van der Meer, F.D. and De Jong, S.M. eds. *Imaging spectrometry: basic principles and prospective applications*. Kluwer Academic Publishers, Dordrecht, 111-155. Remote Sensing and Digital Image Processing no. 4.
- Kumar, L. and Skidmore, A.K., 2000. Radiation-vegetation relationships in a Eucalyptus forest. *Photogrammetric Engineering and Remote Sensing*, 66 (2), 193-204.
- Kunkel, K. and Pletscher, D.H., 2001. Winter hunting patterns of wolves in and near glacier National Park, Montana. *Journal of Wildlife Management*, 65 (3), 520-530.
- Kyriazakis, I. and Oldham, J.D., 1993. Diet selection in sheep: the ability of growing lambs to select a diet that meets their crude protein (nitrogen x 6.25) requirements. *British Journal of Nutrition*, 69 (3), 617-629.
- Kyriazakis, I. and Oldham, J.D., 1997. Food intake and diet selection in sheep: the effect of manipulating the rates of digestion of carbohydrates and protein of the foods offered as a choice. *British Journal of Nutrition*, 77 (2), 243-254.
- Kyriazakis, I., Oldham, J.D., Coop, R.L., et al., 1994. The effect of subclinical intestinal nematode infection on the diet selection of growing sheep. *British Journal of Nutrition*, 72 (5), 665-677.
- Laca, E.A., 1998. Spatial memory and food searching mechanisms of cattle. *Journal of Range Management*, 51 (4), 370-378.

- Laca, E.A. and Demment, M.W., 1996. Foraging strategies of grazing animals. *In: Hodgson, J. and Illius, A.W. eds. The ecology and management of grazing systems.* CAB international, Wallingford, 137-157.
- Laca, E.A., Distel, R.A., Griggs, T.C., et al., 1994a. Effects of canopy structure on patch depression by grazers. *Ecology*, 75 (3), 706-716.
- Laca, E.A., Distel, R.A., Griggs, T.C., et al., 1993. Field test of optimal foraging with cattle: the marginal value theorem successfully predicts patch selection and utilization. *In: Baker, M.J. ed. Grasslands for our world: proceedings of the XVII international grassland congress, held in February 1993, in New Zealand and Queensland.* SIR, Wellington, 709-710.
- Laca, E.A., Ungar, E.D. and Demment, M.W., 1994b. Mechanisms of handling time and intake rate of a large mammalian grazer. *Applied Animal Behaviour Science*, 39 (1), 3-19.
- Laca, E.A., Ungar, E.D., Seligman, N., et al., 1992. Effects of sward height and bulk density on bite dimensions of cattle grazing homogeneous swards. *Grass and Forage Science*, 47 (1), 91-102.
- Lamprey, H.F., 1963. Ecological separation of the large mammal species in the Tarangire Game Reserve, Tanganyika. *East African Wildlife Journal*, 1, 63-92.
- Lane, C. and Moorehead, R., 1994. New directions in rangeland resource tenure and policy. *In: Scoones, I. ed. Living with uncertainty: new directions in pastoral development in Africa.* Intermediate Technology Publications, London, 96-133.
- Langer, P., 1988. *Mammalian herbivore stomach: comparative anatomy, function and evolution.* Fischer, Stuttgart.
- Langvatn, R. and Hanley, T.A., 1993. Feeding-patch choice by red deer in relation to foraging efficiency: an experiment. *Oecologia*, 95 (2), 164-170.
- Lawler, I.R., Foley, W.J., Eschler, B.M., et al., 1998. Intraspecific variation in Eucalyptus secondary metabolites determines food intake by folivorous marsupials. *Oecologia*, 116 (1/2), 160-169.
- Laws, R.M., 1970. Elephants as agents of habitat and landscape change in East Africa. *Oikos*, 21 (1), 1-15.
- Lawson, A., Biggeri, A. and Boehning, D., 1999. *Disease mapping and risk assessment for public health.* Wiley, Chichester.
- Lazo, A., 1994. Social segregation and the maintenance of social stability in a feral cattle population. *Animal Behaviour*, 48 (5), 1133-1141.
- Le Houerou, H.N., 1989. *The grazing land ecosystems of the African Sahel.* Springer, Berlin.
- Ledger, H.P., 1968. Body composition as a basis for a comparative study of some East African mammals. *In: Crawford, M.A. ed. Comparative nutrition of wild animals: proceedings of a symposium held at the Zoological Society of London, on 10 and 11 November 1966.* Academic Press, London, 289-310.
- LeDoux, J.E., 1992. Brain mechanisms of emotion and emotional learning. *Current Opinion in Neurobiology*, 2 (2), 191-197.
- LeDoux, J.E., 1994. Emotion, memory and the brain. *Scientific American*, 270 (6), 50-57.
- Legendre, P. and Fortin, M.J., 1989. Spatial pattern and ecological analysis. *Vegetatio*, 80 (2), 107-138.
- Leuthold, B.M. and Leuthold, W., 1978. Ecology of the giraffe in Tsavo East National Park, Kenya. *East African Wildlife Journal*, 16 (1), 1-20.
- Levin, S.A., 1992. The problem of pattern and scale in ecology: the Robert H. MacArthur Award Lecture. *Ecology*, 73 (6), 1943-1967.
- Levins, R., 1968. *Evolution in changing environments: some theoretical explorations.* Princeton University Press, Princeton.
- Lewis, J.E., Rowland, J. and Nadeau, A., 1998. Estimating maize production in Kenya using NDVI: some statistical considerations. *International Journal of Remote Sensing*, 19 (13), 2609-2617.
- Lewis, J.G., 1977. Game domestication for animal production in Kenya: activity patterns of eland, oryx, buffalo and zebu cattle. *Journal of Agricultural Science, UK*, 89 (3), 551-563.
- Lewis, M.M., 1994. Species composition related to spectral classification in an Australian spinifex hummock grassland. *International Journal of Remote Sensing*, 15 (16), 3223-3239.
- Li, B.L., Gorshkov, V.G. and Makarieva, A.M., 2004. Energy partitioning between different-sized organisms and ecosystem stability. *Ecology*, 85 (7), 1811-1813.
- Li, W., Wang, Z., Ma, Z., et al., 1997. A regression model for the spatial distribution of red-crown crane in Yancheng Biosphere Reserve, China. *Ecological Modelling*, 103 (2), 115-121.
- Li, W.K., 2002. Macroecological patterns of phytoplankton in the northwestern North Atlantic Ocean. *Nature*, 419 (6903), 154-157.
- Lima, S.L., 2002. Putting predators back into behavioral predator-prey interactions. *Trends in Ecology and Evolution*, 17 (2), 70-75.
- Lincoln, D.E., Newton, T.S., Ehrlich, P.R., et al., 1982. Coevolution of the checkerspot butterfly *Euphydryas chalcedona* and its larval food plant *Diplacus aurantiacus*: larval response to protein and leaf resin. *Oecologia*, 52 (2), 216-223.

- Lind, J. and Sheikh, N., 2001. Armaments, environments: small arms and control of natural resources. *Eco-Conflicts*, 1 (3), 1-4. [<http://www.acts.or.ke/pubs/policybriefs/pubs/ecopolicy%20brief2.pdf>]
- Lindsay, R.W., Percival, D.B. and Rothrock, D.A., 1996. The discrete wavelet transform and the scale analysis of the surface properties of sea ice. *IEEE Transactions on Geoscience and Remote Sensing*, 34 (3), 771-787.
- Liu, X., 2001. *Mapping and modelling the habitat of giant pandas in Foping Nature Reserve, China*. Wageningen University, Wageningen. PhD thesis
- Liu, X., Skidmore, A.K., Wang, T., et al., 2002. Giant Panda movements in Foping Nature Reserve, China. *The Journal of Wildlife Management*, 66 (4), 1179-1188.
- Lloyd, L.E., MacDonald, B.E. and Crampton, E.W., 1978. *Fundamentals of nutrition*. Freeman, San Francisco.
- Loeb, S.C. and Hooper, R.G., 1997. An experimental test of interspecific competition for red-cockaded woodpecker cavities. *Journal of Wildlife Management*, 61 (4), 1268-1280.
- Los, S.O., 1998. *Linkages between global vegetation and climate: an analysis based on NOAA-Advanced Very High Resolution Radiometer data*. NASA. NASA Report no. GSFC/CR-1998-206852.
- Loth, P.E. and Prins, H.H.T., 1986. Spatial patterns of the landscape and vegetation of the Lake Manyara National Park. *ITC Journal*, 2, 115-130.
- Low, W.A., Tweedie, R.L., Edwards, C.B.H., et al., 1981. The influence of environment on daily maintenance behaviour of free-ranging Shorthorn cows in central Australia. I. General introduction and descriptive analysis of day-long activities. *Applied Animal Ethology*, 7 (1), 11-26.
- Ludwig, F., 2001. *Tree-grass interactions on an East African savanna: the effects of facilitation, competition, and hydraulic lift*. Wageningen University, Wageningen. PhD thesis
- Ludwig, F., De Kroon, H., Prins, H.H.T., et al., 2001. Effects of nutrients and shade on tree-grass interactions in an East African savanna. *Journal of Vegetation Science*, 12 (4), 579-588.
- MCloskey, R.T., 1980. Spatial patterns in sizes of seeds collected by four species of heteromyid rodents. *Ecology*, 61 (3), 486-489.
- Macandza, V.A., Owen-Smith, N. and Cross, P.C., 2004. Forage selection by African buffalo (*Syncerus caffer*) through the dry season in two landscapes of the Kruger National Park. *South African Journal of Wildlife Research*, 34, 113-121.
- MacArthur, R.H., 1958. Population ecology of some warblers of northeastern coniferous forests. *Ecology*, 39 (4), 599-619.
- MacFadden, B.J. and Shockey, B.J., 1997. Ancient feeding ecology and niche differentiation of Pleistocene mammalian herbivores from Tarija, Bolivia: morphological and isotopic evidence. *Paleobiology*, 23 (1), 77-100.
- Macfadyan, W.A., 1950. Soil and vegetation in British Somaliland. *Nature*, 165 (4186), 121.
- Makkar, H.P.S., 2003. Effects and fate of tannins in ruminant animals, adaptation to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Ruminant Research*, 49 (3), 241-256.
- Mangan, J.L., 1988. Nutritional effects of tannins in animal feeds. *Nutrition Research Reviews*, 1 (1), 209-231.
- Mangel, M. and Clark, C.W., 1988. *Dynamic modelling in behavioral ecology*. Princeton University Press, Princeton.
- Manly, B.F.J., 1999. *Randomization, bootstrap and Monte Carlo methods in biology*. 2nd edn. Chapman & Hall, London.
- Manseau, M., 1996. *Relation réciproque entre les caribous et la végétation des aires d'estivage: le cas du troupeau de caribous de la rivière George*. Université Laval, Laval. PhD thesis
- Marell, A., Ball, J.P. and Hofgaard, A., 2002. Foraging and movement paths of female reindeer: insights from fractal analysis, correlated random walks, and Levy flights. *Canadian Journal of Zoology*, 80 (5), 854-865.
- Markus, S., Bailey, D.W., Jensen, D., et al., 1998. A preliminary evaluation of a fenceless livestock control system. *Journal of Animal Science*, 76 (Suppl. 1), 103.
- Marotti, D.M., Cosgrove, G.P., Chapman, D.F., et al., 2002. Growing grass and clover separately allows animals to sustain a high nutrient intake. *Proceedings of the New Zealand Society of Animal Production*, 62, 163-166.
- Marschner, H., 1995. *Mineral nutrition of higher plants*. 2nd edn. Academic Press, London.
- Martin, M.E., Newman, S.D., Aber, J.D., et al., 1998. Determining forest species composition using high spectral resolution remote sensing data. *Remote Sensing of Environment*, 65 (3), 249-254.
- Martin, P.S., 1982. The pattern and meaning of holarctic mammoth extinction. In: Hopkins, D.M., Matthews, J.V., Schweger, C.S., et al. eds. *The paleoecology of Beringia*. Academic Press, New York, 398-408.

- Martin, P.S. and Wright, H.E., 1967. *Pleistocene extinctions: the search for a cause*. Yale University Press, New Haven.
- Mason, S.J., Waylen, P.R., Mimmack, G.M., et al., 1999. Changes in extreme rainfall events in South Africa. *Climatic Change*, 41 (2), 249-257.
- Matson, K.D., Millam, J.R. and Klasing, K.C., 2004. Cockatiels (*Nymphicus hollandicus*) reject very low levels of plant secondary compounds. *Applied Animal Behaviour Science*, 85 (1/2), 141-156.
- Mautz, W.W., Kanter, J. and Pekins, P.J., 1992. Seasonal metabolic rhythms of captive female white-tailed deer: a reexamination. *The Journal of Wildlife Management*, 56 (4), 656-661.
- Mbogo, S., 2003. Pastoral shoot-outs: breaking the link between small arms and environmental conflict. *Toward Freedom* (51). [<http://towardfreedom.com/home/content/view/106/0/>]
- McCabe, J.T., 1994. Mobility and land use among African pastoralists: old conceptual problems and new interpretations. In: Fratkin, E., Galvin, K.A. and Roth, E.A. eds. *African pastoral systems: an integrated approach*. Lynne Rienner Publishers, Boulder, 69-89.
- McComb, K., Moss, C., Durant, S.M., et al., 2001. Matriarchs as repositories of social knowledge in African elephants. *Science* (292), 491-494.
- McIlvain, E.H. and Shoop, M.C., 1971. Shade for improving cattle gains and rangeland use. *Journal of Range Management*, 24 (3), 181-184.
- McNair, J.N., 1982. Optimal giving-up times and the marginal value theorem. *The American Naturalist*, 119 (4), 511-529.
- McNamara, J.M., 1987. Starvation and predation as factors limiting population size. *Ecology*, 68 (5), 1515-1519.
- McNamara, J.M. and Houston, A.I., 1987. Partial preferences and foraging. *Animal behaviour*, 35 (4), 1084-1099.
- McNaughton, S.J., 1976. Serengeti migratory wildebeest: facilitation of energy flow by grazing. *Science, USA*, 191 (4222), 92-94.
- McNaughton, S.J., 1984. Grazing lawns: animals in herds, plant form, and coevolution. *The American Naturalist*, 124 (6), 863-886.
- McNaughton, S.J., 1985. Ecology of a grazing ecosystem: the Serengeti. *Ecological Monographs*, 55 (3), 259-294.
- McNaughton, S.J., 1990. Mineral nutrition and seasonal movements of African migratory ungulates. *Nature London*, 345 (6276), 613-615.
- McNaughton, S.J., 1995. Plant communities and herbivory. In: Sinclair, A.R.E. and Arcese, P. eds. *Serengeti 2: dynamics, management, and conservation of an ecosystem: based on papers at a workshop held in December 1991 at the Serengeti Research Institute, Tanzania*. University of Chicago Press, Chicago.
- McSweeney, C.S., Palmer, B., McNeill, D.M., et al., 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Animal Feed Science and Technology*, 91 (1), 83-93.
- Mearns, R. and Swift, J., 1996. Pasture tenure and management in the retreat from a centrally planned economy in Mongolia. In: West, N.E. ed. *Rangelands in a sustainable biosphere: proceedings of the fifth international rangeland congress, held in Salt Lake City, Utah, July 23-28, 1995*. Society for Range Management, Denver, 96-98.
- Meissner, H.H. and Paulsmeier, D.V., 1995. Plant compositional constituents affecting between-plant and animal species prediction of forage intake. *Journal of Animal Science*, 73 (8), 2447-2457.
- Mendl, M. and Held, S., 2001. Living in groups: an evolutionary approach. In: Keeling, L.J. and Gonyou, H.W. eds. *Social behaviour in farm animals*. Cabi, Wallingford, 7-31.
- Mendoza, M., Janis, C.M. and Palmqvist, P., 2002. Characterizing complex craniodental patterns related to feeding behaviour in ungulates: a multivariate approach. *Journal of Zoology*, 258 (2), 223-246.
- Mills, M.G.L., 1984. Prey selection and feeding habits of the large carnivores in the southern Kalahari. *Koedoe*, 27 (Suppl.), 281-294.
- Milne, B.T., Turner, M.G., Wiens, J.A., et al., 1992. Interactions between the fractal geometry of landscapes and allometric herbivory. *Theoretical Population Biology*, 41 (3), 337-353.
- Milne, J.A., Hodgson, J., Thompson, R., et al., 1982. The diet ingested by sheep grazing swards differing in white clover and perennial ryegrass content. *Grass and Forage Science*, 37 (3), 209-218.
- Mishra, C., Prins, H.H.T. and Van Wieren, S.E., 2001. Overstocking in the trans-Himalayan rangelands of India. *Environmental Conservation*, 28 (3), 279-283.
- Mishra, C., Prins, H.H.T. and Van Wieren, S.E., 2003. Diversity, risk mediation, and change in a trans-Himalayan agropastoral system. *Human Ecology*, 31 (4), 595-609.
- Mishra, C., Van Wieren, S.E., Ketner, P., et al., 2004. Competition between domestic livestock and wild bharal *Pseudois nayaur* in the Indian Trans-Himalaya. *Journal of Applied Ecology*, 41 (2), 344-354.
- Moen, A.N., 1973. *Wildlife ecology: an analytical approach*. Freeman, San Francisco.

- Moen, R., 1998. Linking moose population and plant growth models with a moose energetics model. *Ecosystems*, 1 (1), 52-63.
- Moen, R., Pastor, J. and Cohen, Y., 1997. A spatially explicit model of moose foraging and energetics. *Ecology*, 78 (2), 505-521.
- Mole, S., 1989. Polyphenolics and the nutritional ecology of herbivores. In: Cheeke, P.R. ed. *Toxicants of plant origin. IV. Phenolics*. CRC Press, Boca Raton, 191-221.
- Monteith, J.L., 1972. Solar radiation and productivity in tropical ecosystems. *Journal of Applied Ecology*, 9 (3), 747-766.
- Monteith, J.L., 1977. Climate and the efficiency of crop production in Britain. *Philosophical Transactions of the Royal Society of London, B*, 281, 277-294.
- Mooney, H.A., 1986. Photosynthesis. In: Crawley, M.J. ed. *Plant ecology*. Blackwell, Oxford, 345-373.
- Moore, D.S., 2002. *The dependent gene: the fallacy of nature vs. nurture*. Henry Holt and Company, New York.
- Moreau, S., Bosseno, R., Gu, X.F., et al., 2003. Assessing the biomass dynamics of Andean bofedal and totora high-protein wetland grasses from NOAA/AVHRR. *Remote Sensing of Environment*, 85 (4), 516-529.
- Morin, P.J., 2000. Biodiversity's ups and downs. *Nature*, 406 (6795), 508-512.
- Moritz, M., Scholte, P. and Kari, S., 2002. The demise of the nomadic contract: arrangements and rangelands under pressure in the far north of Cameroon (1). *Nomadic Peoples*, 6 (1), 124-144.
- Morris, D.W., Fox, B.J., Luo, J., et al., 2000. Habitat-dependent competition and the coexistence of Australian heathland rodents. *Oikos*, 91 (2), 294-306.
- Morrison, M.L. and Hall, L.S., 2002. Standard terminology: toward a common language to advance ecological understanding and application. In: Scott, J.M. ed. *Predicting species occurrences: issues of accuracy and scale*. Island Press, Washington.
- Morse, D.R., Lawton, J.H., Dodson, M.M., et al., 1985. Fractal dimension of vegetation and the distribution of arthropod body lengths. *Nature, UK*, 314 (6013), 731-733.
- Mott MacDonald, 1999. *Logone Floodplain model study report*. Mott MacDonald Group, Cambridge.
- Mueggler, W.F., 1965. Cattle distribution on steep slopes. *Journal of Range Management*, 18 (5), 255-257.
- Murray, M.G., 1995. Specific nutrient requirements and migration of wildebeest. In: Sinclair, A.R.E. and Arcese, P. eds. *Serengeti 2: dynamics, management, and conservation of an ecosystem: based on papers at a workshop held in December 1991 at the Serengeti Research Institute, Tanzania*. University of Chicago Press, Chicago, 231-256.
- Murray, M.G., 2003. Overkill and sustainable use. *Science*, 299 (5614), 1851 - 1853.
- Murray, M.G. and Brown, D., 1993. Niche separation of grazing ungulates in the Serengeti: an experimental test. *The Journal of Animal Ecology*, 62 (2), 380-389.
- Murwira, A., 2003. *Scale matters!: a new approach to quantify spatial heterogeneity for predicting the distribution of wildlife*. Wageningen University, Wageningen. PhD thesis [<http://www.library.wur.nl/wda/dissertations/dis3506.pdf>]
- Mutanga, O. and Skidmore, A.K., 2004a. Integrating imaging spectroscopy and neural networks to map grass quality in the Kruger National Park, South Africa. *Remote Sensing of Environment*, 90 (1), 104-115.
- Mutanga, O. and Skidmore, A.K., 2004b. Narrow band vegetation indices overcome the saturation problem in biomass estimation. *International Journal of Remote Sensing*, 25 (19), 3999-4014.
- Mutanga, O., Skidmore, A.K. and Prins, H.H.T., 2004a. Discriminating sodium concentration in a mixed grass species environment of the Kruger National Park using field spectrometry. *International Journal of Remote Sensing*, 25 (20), 4191-4201.
- Mutanga, O., Skidmore, A.K. and Prins, H.H.T., 2004b. Predicting in situ pasture quality in the Kruger National Park, South Africa, using continuum-removed absorption features. *Remote Sensing of Environment*, 89 (3), 393-408.
- Mwasi, S.M., 2002. *Compressed nature: co-existing grazers in a small reserve in Kenya*. Wageningen University, Wageningen. PhD thesis
- Mysterud, A., Lian, L.B. and Hjernann, D.O., 1999. Scale-dependent trade-offs in foraging by European roe deer (*Capreolus capreolus*) during winter. *Canadian Journal of Zoology*, 77 (9), 1486-1493.
- Mysterud, A., Yoccoz, N.G., Stenseth, N.C., et al., 2001. Effects of age, sex and density on body weight of Norwegian red deer: evidence of density-dependent senescence. *Proceedings of the Royal Society of London Series B, Biological Sciences*, 268 (1470), 911-919.
- Naah, E., 1992. *Hydrologie du Grand Yaéré du nord-Cameroun*. Université de Yaoundé, Yaoundé. PhD thesis

- Nagy, K.A., Girard, I.A. and Brown, T.K., 1999. Energetics of free-ranging mammals, reptiles, and birds. *Annual Review of Nutrition*, 19, 247-277.
- NCAA, 2000. *1998 aerial boma count, 1999 people and livestock census, and human population trend between 1954 and 1999 in the NCA*. Ngorongoro Conservation Area Authority, Ngorongoro Crater.
- Nellemann, C., 1997. Grazing strategies of muskoxen (*Ovibos moschatus*) during winter in Angujaartorfiup Nunaa in western Greenland. *Canadian Journal of Zoology*, 75 (7), 1129-1134.
- Newman, J.A., Parsons, A.J. and Harvey, A., 1992. Not all sheep prefer clover: diet selection revisited. *Journal of Agricultural Science*, 119 (2), 275-283.
- Newman, J.A., Parsons, A.J., Thornley, J.H.M., et al., 1995. Optimal diet selection by a generalist grazing herbivore. *Functional Ecology*, 9 (2), 255-268.
- Niamir-Fuller, M., 1999. *Managing mobility in African Rangelands: the legitimization of transhumance*. Intermediate Technology Publications, London.
- Niamir-Fuller, M. and Turner, M.D., 1999. A review of recent literature on pastoralism and transhumance in Africa. In: Niamir-Fuller, M. ed. *Managing mobility in African Rangelands: the legitimization of transhumance*. Intermediate Technology Publications, London, 18-46.
- Northup, R.R., Yu, Z.S., Dahlgren, R.A., et al., 1995. Polyphenol control of nitrogen release from pine litter. *Nature London*, 377 (6546), 227-229.
- Norton-Griffiths, M., 1975. *Counting animals*. Serengeti Ecological Monitoring Programme, Nairobi.
- Novellie, P.A., 1978. Comparison of the foraging strategies of blesbok and springbok on the Transvaal highveld. *South African Journal of Wildlife Research*, 8, 137-144.
- Novellie, P.A., Manson, J. and Bigalke, R.C., 1984. Behavioural ecology and communication in the Cape grysbok. *South African Journal of Zoology*, 19 (1), 22-30.
- O'Neill, R.V., Gardner, R.H., Milne, B.T., et al., 1991. Heterogeneity and spatial hierarchies. In: Kolasa, J., Pickett, S.T.A., Allen, T.F.H., et al. eds. *Ecological heterogeneity*. Springer, New York, 85-96.
- O'Reagain, P.J., Goetsch, B.C. and Owen-Smith, R.N., 1996. Effect of species composition and sward structure on the ingestive behaviour of cattle and sheep grazing South African sourveld. *Journal of Agricultural Science*, 127 (2), 271-280.
- Oates, J.F., Waterman, P.G. and Choo, G.M., 1980. Food selection by the South Indian leaf-monkey, *Presbytis johnii*, in relation to leaf chemistry. *Oecologia*, 45 (1), 45-56.
- Oosterheld, M., DiBella, C.M. and Kerdiles, H., 1998. Relation between NOAA-AVHRR satellite data and stocking rate of rangelands. *Ecological Applications*, 8 (1), 207-212.
- Oosterheld, M., Sala, O.E. and McNaughton, S.J., 1992. Effect of animal husbandry on herbivore-carrying capacity at a regional scale. *Nature*, 356 (6366), 234-236.
- Oftedal, O.T., 1984. Milk composition, milk yield and energy output at peak lactation: a comparative review. *Symposia of the Zoological Society of London*, 51 (Physiological strategies in lactation), 33-85.
- Oftedal, O.T., 1985. Pregnancy and lactation. In: Hudson, R.J. and White, R.G. eds. *Bioenergetics of wild herbivores*. CRC, Boca Raton, 215-238.
- Ogden, R.T., 1997. *Essential wavelets for statistical applications and data analysis*. Birkhaeuser, Boston.
- Ogura, S., Hasegawa, H. and Hirata, M., 2002. Effects of herbage mass and herbage quality on spatially heterogeneous grazing by cattle in a bahia grass (*Paspalum notatum*) pasture. *Tropical Grasslands*, 36 (3), 172-179.
- Oindo, B.O., 2001. *Spatial patterns of species diversity in Kenya*. Wageningen University, Wageningen. PhD thesis
- Oindo, B.O., 2002. Predicting mammal species richness and abundance using multi-temporal NDVI. *PEandRS, Photogrammetric Engineering and Remote Sensing*, 68 (6), 623-629.
- Oindo, B.O., De By, R.A. and Skidmore, A.K., 2000. Interannual variability of NDVI and bird species diversity in Kenya. *International Journal of Applied Earth Observation and Geoinformation*, 2 (3), 172-180.
- Oindo, B.O., Skidmore, A.K. and Prins, H.H.T., 2001. Body size and abundance relationship: an index of diversity for herbivores. *Biodiversity and Conservation*, 10 (11), 1923-1931.
- Ole Katampoi, K., Genga, G.O., Mwangi, M., et al., 1990. *Kajiado District atlas*. Arid and Semi Arid Lands (ASAL) Programme, Kajiado.
- Oloff, H., De Leeuw, J., Bakker, J.P., et al., 1997. Vegetation succession and herbivory in a salt marsh: changes induced by sea level rise and silt deposition along an elevational gradient. *Journal of Ecology Oxford*, 85 (6), 799-814.
- Oloff, H., Ritchie, M.E. and Prins, H.H.T., 2002. Global environmental determinants of diversity in large herbivores. *Nature*, 415, 901-905.
- Olivry, J.C., Chouret, A., Vuillaume, G., et al., 1996. *Hydrologie du Lac Tchad*. ORSTOM editions, Paris.

- Olton, D.S., 1978. Characteristics of spatial memory. *In: Hulse, S.H., Fowler, H. and Honig, W.K. eds. Cognitive processes in animal behavior.* Erlbaum, Hillsdale, 341-373.
- Orlove, B.S., 1977. Integration through production: the use of zonation in Espinar. *American Ethnologist*, 4 (1), 84-101.
- Ottichilo, W.K., 2000. *Wildlife dynamics: an analysis of change in the Masai Mara ecosystem of Kenya.* Wageningen University, Wageningen. PhD thesis
- Ottichilo, W.K., De Leeuw, J. and Prins, H.H.T., 2001. Population trends of resident wildebeest [*Connochaetes taurinus hecki* (Neumann)] and factors influencing them in the Masai Mara ecosystem, Kenya. *Biological Conservation*, 97 (3), 271-282.
- Ottichilo, W.K., De Leeuw, J., Skidmore, A.K., et al., 2000a. Population trends of large non-migratory wild herbivores and livestock in the Masai Mara ecosystem, Kenya, between 1977 and 1997. *African Journal of Ecology*, 38 (3), 202-216.
- Ottichilo, W.K., Grunblatt, J., Said, M.Y., et al., 2000b. Wildlife and livestock population trends in the Kenya rangeland. *In: Prins, H.H.T., Grootenhuis, J.G. and Dolan, T.T. eds. Wildlife conservation by sustainable use.* Kluwer Academic Publishers, Boston, 203-218.
- Owen-Smith, N., 1975. The social ethology of the white rhinoceros. *Zeitschrift für Tierpsychologie*, 38, 337-384.
- Owen-Smith, N., 1982. Factors influencing the consumption of plant products by large herbivores. *In: Huntley, B.J. and Walker, B.H. eds. The ecology of tropical savannas.* Springer-Verlag, Berlin, 359-404.
- Owen-Smith, N., 1989. Morphological factors and their consequences for resource partitioning among African savanna ungulates: a simulation modelling approach. *In: Morris, D.W., Abramsky, Z., Fox, B.J., et al. eds. Patterns in the structure of mammalian communities.* Texas Tech University Press, Lubbock, 155-165.
- Owen-Smith, N., 1993. Evaluating optimal diet models for an African browsing ruminant, the kudu: how constraining are the assumed constraints? *Evolutionary Ecology*, 7 (5), 499-524.
- Owen-Smith, N., 1994. Foraging responses of kudus to seasonal changes in food resources: elasticity in constraints. *Ecology*, 75 (4), 1050-1062.
- Owen-Smith, N., 1996. Circularity in linear programming models of optimal diet. *Oecologia*, 108 (2), 259-261.
- Owen-Smith, N., 1998. How high ambient temperature affects the daily activity and foraging time of a subtropical ungulate, the greater kudu (*Tragelaphus strepsiceros*). *Journal of Zoology*, 246 (02), 183-192.
- Owen-Smith, N., 2000. Modeling the population dynamics of a subtropical ungulate in a variable environment: rain, cold and predators. *Natural Resource Modeling*, 13, 57-87.
- Owen-Smith, N., 2002a. *Adaptive herbivore ecology: from resources to populations in variable environments.* Cambridge University Press, Cambridge.
- Owen-Smith, N., 2002b. Credible models for herbivore-vegetation systems: towards an ecology of equations. *South African Journal of Science*, 98 (9/10), 445-449.
- Owen-Smith, N., 2002c. A metaphysiological modelling approach to stability in herbivore-vegetation systems. *Ecological Modelling*, 149 (1), 153-178.
- Owen-Smith, N., 2004. Functional heterogeneity in resources within landscapes and herbivore population dynamics. *Landscape Ecology*, 19 (7), 761-771.
- Owen-Smith, N. and Cooper, S.M., 1987. Palatability of woody plants to browsing ruminants in a South African savanna. *Ecology, USA*, 68 (2), 319-331.
- Owen-Smith, N. and Cooper, S.M., 1989. Nutritional ecology of a browsing ruminant, the kudu (*Tragelaphus strepsiceros*), through the seasonal cycle. *Journal of Zoology*, 219 (1), 29-43.
- Owen-Smith, N. and Cumming, D.H.M., 1993. Comparative foraging strategies of grazing ungulates in African savanna grasslands. *In: Baker, M. ed. Grasslands for our world: proceedings of the 17th International Grassland Congress* SIR Publishing, 691-697.
- Owen-Smith, N. and Danckwerts, J.E., 1997. Herbivory. *In: Cowling, R.M., Richardson, D.M. and Pierce, S.M. eds. Vegetation of Southern Africa.* Cambridge University Press, Cambridge.
- Owen-Smith, N. and Novellie, P., 1982. What should a clever ungulate eat? *The American Naturalist*, 119 (2), 151-178.
- Owen-Smith, R.N., 1985. Niche separation among African ungulates. *In: Vrba, E.S. ed. Species and speciation.* Transvaal Museum, Pretoria, 161-171. Transvaal Museum Monograph No. 4.
- Owen-Smith, R.N., 1988. *Megaherbivores: the influence of very large body size on ecology.* Cambridge University Press, Cambridge.

- Owens, M.K., Launchbaugh, K.L. and Holloway, J.W., 1991. Pasture characteristics affecting spatial distribution of utilization by cattle in mixed brush communities. *Journal of Range Management*, 44 (2), 118-123.
- Oxby, C., 1982. *Group ranches in Africa*. Overseas Development Institute, London. Pastoral Network Paper Agricultural Administration Unit, Overseas Development Institute.
- Paine, R.T., 1969. A note on trophic complexity and community stability. *The American Naturalist*, 103 (929), 91-93.
- Palmer, M.W., 1992. The coexistence of species in fractal landscapes. *The American Naturalist*, 139 (2), 375-397.
- Parker, I.S.C., 1983. The Tsavo story: an ecological case history. In: Owen-Smith, R.N. ed. *Management of large mammals in African conservation areas: proceedings of a symposium, held in Pretoria, South Africa, 29-30 April 1982*. Haum, Pretoria, 37-50.
- Parker, K.L., 1988. Effects of heat, cold, and rain on coastal black-tailed deer. *Canadian Journal of Zoology*, 66 (11), 2475-2483.
- Parker, K.L., Gillingham, M.P., Hanley, T.A., et al., 1993. Seasonal patterns in body mass, body composition, and water transfer rates of free-ranging and captive black-tailed deer (*Odocoileus hemionus sitkensis*) in Alaska. *Canadian Journal of Zoology*, 71, 1397-1404.
- Parker, K.L., Gillingham, M.P., Hanley, T.A., et al., 1999. *Energy and protein balance of free-ranging black-tailed deer in a natural forest environment*. Wildlife Monographs no. 143.
- Parlar, M. and Weng, Z.K., 2003. Balancing desirable but conflicting objectives in the newsvendor problem. *IIE Transactions*, 35 (2), 131-142.
- Parsons, A.J., Newman, J.A., Penning, P.D., et al., 1994a. Diet preference of sheep: effects of recent diet, physiological state and species abundance. *Journal of Animal Ecology*, 63 (2), 465-478.
- Parsons, A.J., Thornley, J.H.M., Newman, J., et al., 1994b. A mechanistic model of some physical determinants of intake rate and diet selection in a two-species temperate grassland sward. *Functional Ecology*, 8 (2), 187-204.
- Paruelo, J.M., Epstein, H.E., Lauenroth, W.K., et al., 1997. ANPP estimates from NDVI for the central grassland region of the United States. *Ecology*, 78 (3), 953-958.
- Perez, C., Ackroff, K. and Sclafani, A., 1996. Carbohydrate-and protein-conditioned flavor preferences: effects of nutrient preloads. *Physiology and Behavior*, 59 (3), 467-474.
- Peters, R.H., 1983. *The ecological implications of body size*. Cambridge University Press, Cambridge.
- Petren, K. and Case, T.J., 1996. An experimental demonstration of exploitation competition in an ongoing invasion. *Ecology*, 77 (1), 118-132.
- Petren, K. and Case, T.J., 1998. Habitat structure determines competition intensity and invasion success in gecko lizards. *Proceedings National Academy of Sciences USA*, 95 (20), 11739-11744.
- Pfister, J.A., Provenza, F.D., Manners, G.D., et al., 1997. Tall larkspur ingestion: can cattle regulate intake below toxic levels? *Journal of Chemical Ecology*, 23 (3), 759-777.
- Pickup, G. and Smith, D.M.S., 1993. Problems, prospects and procedures for assessing the sustainability of pastoral land management in arid Australia. *Journal of Biogeography*, 20 (5), 471-487.
- Pielou, E.C., 1984. *The interpretation of ecological data: a primer on classification and ordination*. Wiley, New York.
- Pinchak, W.E., Smith, M.A., Hart, R.H., et al., 1991. Beef cattle distribution patterns on foothill range. *Journal of Range Management*, 41 (3), 267-275.
- Pitt, W.C. and Ritchie, M.E., 2002. Influence of prey distribution on the functional response of lizards. *Oikos*, 96 (1), 157-163.
- Pollard, J.C. and Littlejohn, R.P., 1996. The effects of pen size on the behaviour of farmed red deer stags confined in yards. *Applied Animal Behaviour Science*, 47 (3/4), 247-253.
- Price, E.O., 1978. Genotype versus experience effects on aggression in wild and domestic Norway rats. *Behaviour* 64 (3/4), 340-353.
- Price, E.O., 1984. Behavioral aspects of animal domestication. *Quarterly Review of Biology*, 59 (1), 1-32.
- Prins, H.H.T., 1987a. *The buffalo of Manyara: the individual in the context of herd life in a seasonal environment of East Africa*. University of Groningen, Groningen. PhD thesis
- Prins, H.H.T., 1987b. Nature conservation as an integral part of optimal land use in East Africa: the case of the Masai Ecosystem of northern Tanzania. *Biological Conservation*, 40 (2), 141-161.
- Prins, H.H.T., 1988. Plant phenology patterns in Lake Manyara National Park, Tanzania. *Journal of Biogeography*, 15 (3), 465-480.
- Prins, H.H.T., 1989a. Buffalo herd structure and its repercussions for condition of individual African buffalo cows. *Ethology*, 81 (1), 47-71.
- Prins, H.H.T., 1989b. Condition changes and choice of social environment in African buffalo bulls. *Behaviour*, 108 (3), 297-324.

- Prins, H.H.T., 1992. The pastoral road to extinction: competition between wildlife and traditional pastoralism in East Africa. *Environmental Conservation*, 19 (2), 117-123.
- Prins, H.H.T., 1996. *Ecology and behaviour of the African buffalo: social inequality and decision making*. Chapman & Hall, London.
- Prins, H.H.T., 1998. The origins of grassland communities in northwestern Europe. In: Wallis De Vries, M.F., Bakker, J.P. and Van Wieren, S.E. eds. *Grazing and conservation management*. Kluwer, Dordrecht, 55-105.
- Prins, H.H.T., 1999. Comment on Ruttan and Borgerhoff-Mulder. *Current Anthropology*, 40, 643-645.
- Prins, H.H.T., 2000. Competition between wildlife and livestock. In: Prins, H.H.T., Grootenhuis, J.G. and Dolan, T.T. eds. *Wildlife conservation by sustainable use*. Kluwer Academic Publishers, Boston, 51-80.
- Prins, H.H.T. and Beekman, J.H., 1989. A balanced diet as a goal for grazing: the food of the Manyara buffalo. *African Journal of Ecology*, 27 (3), 241-259.
- Prins, H.H.T. and Douglas-Hamilton, I., 1990. Stability in a multi-species assemblage of large herbivores in East Africa. *Oecologia*, 83 (3), 392-400.
- Prins, H.H.T., Grootenhuis, J.G. and Dolan, T.T., 2000. *Wildlife conservation by sustainable use*. Kluwer Academic Publishers, Boston.
- Prins, H.H.T. and Iason, G.R., 1989. Dangerous lions and nonchalant buffalo. *Behaviour*, 108 (3), 262-293.
- Prins, H.H.T. and Loth, P.E., 1988. Rainfall patterns as background to plant phenology in northern Tanzania. *Journal of Biogeography*, 15 (3), 451-463.
- Prins, H.H.T. and Olf, H., 1998. Species-richness of African grazer assemblages: towards a functional explanation. In: Newbery, D.M., Prins, H.H.T. and Brown, N.D. eds. *Dynamics of tropical communities: the 37th symposium of the British Ecological Society, Cambridge University, 1996*. Blackwell Science, Oxford, 449-490.
- Provenza, F.D., 1995. Postingestive feedback as an elementary determinant of food preference and intake in ruminants. *Journal of Range Management*, 48 (1), 2-17.
- Provenza, F.D., 1996. Acquired aversions as the basis for varied diets of ruminants foraging on rangelands. *Journal of Animal Science*, 74 (8), 2010-2020.
- Provenza, F.D., 2003. Twenty-five years of paradox in plant-herbivore interactions and "sustainable" grazing management. *Rangelands*, 25 (6), 4-15.
- Provenza, F.D., Lynch, J.J. and Nolan, J.V., 1994. Food aversion conditioned in anesthetized sheep. *Physiology and Behavior*, 55 (3), 429-432.
- Provenza, F.D. and Malechek, J.C., 1984. Diet selection by domestic goats in relation to blackbrush twig chemistry. *Journal of Applied Ecology*, 21, 831-841.
- Provenza, F.D., Pfister, J.A. and Cheney, C.D., 1992. Mechanisms of learning in diet selection with reference to phytotoxicosis in herbivores. *Journal of Range Management*, 45 (1), 36-45.
- Provenza, F.D., Scott, C.B., Phy, T.S., et al., 1996. Preference of sheep for foods varying in flavors and nutrients. *Journal of Animal Science*, 74, 2355-2361.
- Provenza, F.D., Villalba, J.J., Dziba, L.E., et al., 2003. Linking herbivore experience, varied diets, and plant biochemical diversity. *Small Ruminant Research*, 49 (3), 257-274.
- Ralphs, M.H. and Provenza, F.D., 1999. Conditioned food aversions: principles and practices, with special reference to social facilitation. *Proceedings of the Nutrition Society*, 58 (4), 813-820.
- Ramirez, I., 1997. Intra-gastric carbohydrate exerts both intake-stimulating and intake-suppressing effects. *Behavioral Neuroscience*, 111 (3), 612-622.
- Redbo, I., 1992. The influence of restraint on the occurrence of oral stereotypies in dairy cows. *Applied Animal Behaviour Science*, 35 (2), 115-123.
- Redbo, I., Redbo-Torstensson, P., Odberg, F.O., et al., 1998. Factors affecting behavioural disturbances in race-horses. *Animal Science*, 66 (2), 475-481.
- Redfern, J.V., Ryan, S.J. and Getz, W.M., 2006. Defining herbivore assemblages in the Kruger National Park: a correlative coherence approach. *Oecologia*, 146 (4), 632-640.
- Regelin, W.L., Schwartz, C.C. and Franzmann, A.W., 1985. Seasonal energy metabolism of adult moose. *Journal of Wildlife Management*, 49 (2), 388-393.
- Reid, R.L., Jung, G.A. and Thayne, W.V., 1988. Relationships between nutritive quality and fiber components of cool season and warm season forages: a retrospective study. *Journal of Animal Science*, 66 (5), 1275.
- Reinhardt, V. and Reinhardt, A., 1981. Cohesive relationships in a cattle herd (*Bos indicus*). *Behaviour*, 77, 121-151.
- Renecker, L.A. and Hudson, R.J., 1989. Seasonal activity budgets of moose in aspen-dominated boreal forests. *The Journal of Wildlife Management*, 53 (2), 296-302.

- Richardson, A.J., Everitt, J.H. and Gausman, H.W., 1983. Radiometric estimation of biomass and nitrogen content of Alicia grass. *Remote Sensing of Environment*, 13 (2), 179-184.
- Ricklefs, R.E. and Schluter, D., 1993. *Species diversity in ecological communities: historical and geographical perspectives*. University of Chicago Press, Chicago.
- Ridout, M.S. and Robson, M.J., 1991. Diet composition of sheep grazing grass/white clover swards: a re-evaluation. *New Zealand Journal of Agricultural Research*, 34 (1), 89-93.
- Rietkerk, M., Boerlijst, M., Van Langevelde, F., et al., 2002a. Self-organization of vegetation in arid ecosystems. *American Naturalist*, 160 (4), 524-530.
- Rietkerk, M., Van de Koppel, J., Kumar, L., et al., 2002b. The ecology of scale. *Ecological Modelling*, 149 (1/2), 1-4.
- Ripley, B.D., 1981. *Spatial statistics*. Wiley, New York.
- Ripple, W.J. and Beschta, R.L., 2003. Wolf reintroduction, predation risk, and cottonwood recovery in Yellowstone National Park. *Forest Ecology and Management*, 184 (1/3), 299-313.
- Ripple, W.J., Larsen, E.J., Renkin, R.A., et al., 2001. Trophic cascades among wolves, elk and aspen on Yellowstone National Park's northern range. *Biological Conservation*, 102 (3), 227-234.
- Risenhoover, K.L., 1986. Winter activity patterns of moose in interior Alaska. *Journal of Wildlife Management*, 50 (4), 727-734.
- Ritchie, M.E., 2002. Competition and coexistence in mobile animals. In: Sommer, U. and Worm, B. eds. *Competition and coexistence*. Springer, Berlin, 127-141.
- Ritchie, M.E. and Olff, H., 1999. Spatial scaling laws yield a synthetic theory of biodiversity. *Nature*, 400 (6744), 557-560.
- Ritchie, M.E. and Tilman, D., 1993. Predictions of species interactions from consumer-resource theory: experimental tests with grasshoppers and plants. *Oecologia*, 94 (4), 516-527.
- Ritchie, M.E., Tilman, D. and Knops, J.M.H., 1998. Herbivore effects on plant and nitrogen dynamics in oak savanna. *Ecology*, 79 (1), 165-177.
- Roath, L.R. and Krueger, W.C., 1982. Cattle grazing and behaviour on a forested range. *Journal of Range Management*, 35 (3), 332-338.
- Robbins, C.T., 1993. *Wildlife feeding and nutrition*. Academic Press, San Diego.
- Robbins, C.T., Mole, S., Hagerman, A.E., et al., 1987. Role of tannins in defending plants against ruminants: reduction in dry matter digestion? *Ecology, USA*, 68 (6), 1606-1615.
- Robbins, C.T., Spalinger, D.E. and Van Hoven, W., 1995. Adaptation of ruminants to browse and grass diets: are anatomical-based browser-grazer interpretations valid? *Oecologia*, 103 (2), 208-213.
- Roberts, S.C. and Dunbar, R.I.M., 1991. Climatic influences on the behavioural ecology of Chanler's mountain reedbuck in Kenya. *African Journal of Ecology*, 29 (4), 316-329.
- Roberts, W.A. and Van Veldhuizen, N., 1985. Spatial memory in pigeons on the radial maze. *Journal of Experimental Psychology. Animal Behavior Processes* 11, 241-260.
- Rosenzweig, M.L., 1968. Net primary productivity of terrestrial communities: prediction from climatological data. *The American Naturalist*, 102 (923), 67-74.
- Rosenzweig, M.L. and Abramsky, Z., 1993. How are diversity and productivity related? In: Ricklefs, R.E. and Schluter, D. eds. *Species diversity in ecological communities: historical and geographical perspectives*. University of Chicago Press, Chicago.
- Rosenzweig, M.L. and Abramsky, Z., 1997. Two gerbils of the Negev: a long-term investigation of optimal habitat selection and its consequences. *Evolutionary Ecology*, 11 (6), 733-756.
- Ruel, J.J. and Ayres, M.P., 1999. Jensen's inequality predicts effects of environmental variation. *Trends in Ecology and Evolution*, 14 (9), 361-366.
- Ruttan, L.M. and Borgerhoff-Mulder, M., 1999. Are East African pastoralists truly conservationists. *Current Anthropology*, 40 (5), 621-652.
- Rutten, M.M.E.M., 1992. *Selling wealth to buy poverty: the process of the individualization of landownership among Maasai pastoralists of Kajiado District, Kenya, 1890 - 1990*. Breitenbach, Fort Lauderdale.
- Ruyle, G.B., Hasson, O. and Rice, R.W., 1987. The influence of residual stems on biting rates of cattle grazing *Eragrostis lehmanniana* Nees. *Applied Animal Behaviour Science*, 19 (1/2), 11-17.
- Said, M.Y., 2003. *Multiscale perspectives of species richness in East Africa*. PhD Thesis, Wageningen University, Wageningen. [<http://www.library.wur.nl/wda/dissertations/dis3351.pdf>]
- Sandford, S., 1994. Improving the efficiency of opportunism: new directions for pastoral development. In: Scoones, I. ed. *Living with uncertainty: new directions in pastoral development in Africa*. Intermediate Technology Publications, London, 174-182.
- Sankhyan, S.K., Bhatta, R., Shinde, A.K., et al., 2001. Seasonal changes in dietary composition and rumen fermentation pattern of sheep on semi-arid rangeland. *Indian Journal of Animal Sciences*, 71 (2), 162-164.

- Savory, A., 1988. *Holistic resource management*. Island Press, Washington.
- Schaefer, J.A. and Messier, F., 1995. Habitat selection as a hierarchy: the spatial scales of winter foraging by muskoxen. *Ecography*, 18 (4), 333-344.
- Schaller, G.B. and Junrang, R., 1988. Effects of a snowstorm on Tibetan antelope. *Journal of Mammalogy*, 69 (3), 631-634.
- Schareika, N., 2001. Environmental knowledge and pastoral migration among the Wodaabe of South-Eastern Niger (1). *Nomadic Peoples*, 5 (1), 65-89.
- Schmidt, K.S. and Skidmore, A.K., 2001. Exploring spectral discrimination of grass species in African rangelands. *International Journal of Remote Sensing*, 22 (17), 3421-3434.
- Schmidt, K.S. and Skidmore, A.K., 2003. Spectral discrimination of vegetation types in a coastal wetland. *Remote Sensing of Environment*, 85 (1), 92-108.
- Schmidt, K.S., Skidmore, A.K., Kloosterman, E.H., et al., 2004. Mapping coastal vegetation using an expert system and hyperspectral imagery. *PEandRS, Photogrammetric Engineering and Remote Sensing*, 70 (6), 703-715.
- Schmitz, O.J., 1991. Thermal constraints and optimization of winter feeding and habitat choice in white-tailed deer. *Holarctic Ecology*, 14 (2), 104-111.
- Schmitz, O.J., Adler, F.R. and Agrawal, A.A., 2003. Linking individual-scale trait plasticity to community dynamics: trait plasticity and community dynamics. *Ecology*, 84 (5), 1081-1082.
- Schmitz, O.J., Cohon, J.L., Rothley, K.D., et al., 1997. Reconciling variability and optimal behaviour using multiple criteria in optimization models. *Evolutionary Ecology*, 12 (1), 73-94.
- Schoener, T.W., 1971. Theory of feeding strategies. *Annual Review of Ecology and Systematics*, 2, 369-404.
- Schoener, T.W., 1973. Population growth regulated by intraspecific competition for energy or time: some simple representations. *Theoretical Population Biology*, 4 (1), 56-84.
- Schoener, T.W., 1976. The species-area relation within archipelagoes: models and evidence from land birds. In: Frith, H.J. and Calaby, J.H. eds. *Proceedings of the 16th International Ornithological Congress*. Australian Academy of Sciences, Canberra, 629-642.
- Schoener, T.W., 1983. Field experiments on interspecific competition. *The American Naturalist*, 122 (2), 240-285.
- Scholes, R.J., 1997. Savanna. In: Cowling, R.M., Richardson, D.M. and Pierce, S.M. eds. *Vegetation of Southern Africa*. Cambridge University Press, Cambridge.
- Scholes, R.J. and Walker, B.H., 1993. *An African savanna: synthesis of the Nylsvley study*. Cambridge University Press, Cambridge.
- Scholte, P., 2005. *Floodplain rehabilitation and the future of conservation & development: adaptive management of success in Waza-Logone, Cameroon*. Wageningen University and Research Centre, Wageningen. Tropical Resource Management Papers no. 67. Ook verschenen als proefschrift Leiden Universiteit, 2005
- Scholte, P., Kari, S., Moritz, M., et al., 2006. Pastoralist responses to floodplain rehabilitation in North Cameroon. *Human Ecology*, 34 (1), 27-51.
- Scholte, P., Kirda, P., Adam, S., et al., 2000. Floodplain rehabilitation in north Cameroon: impact on vegetation dynamics. *Applied Vegetation Science*, 3 (1), 33-42.
- Scholte, P., Pamo, E., Kari, S., et al., 1996. Floodplain rehabilitation in North Cameroon: expected impact on vegetation, pastoralists and wildlife. In: West, N.E. ed. *Rangelands in a sustainable biosphere: proceedings of the fifth international rangeland congress*. Society for Range Management, Denver, 492-493.
- Scholte, P.T., 1992. Leaf litter and Acacia pods as feed for livestock during the dry season in Acacia-Commiphora bushland, Kenya. *Journal of Arid Environments*, 22 (3), 271-276.
- Schrader, T., 1986. *Les Yaérés au Nord du Cameroun: pâturages de saison sèche ?*. CML/Ecole de Faune, Leiden.
- Sclafani, A., 2000. Macronutrient-conditioned flavour preferences. In: Berthoud, H.J. and Seeley, R.J. eds. *Neural and metabolic control of macronutrient intake*. CRC Press, New York, 93-106.
- Scoones, I., 1991. Wetlands in drylands: key resources for agricultural and pastoral production in Africa. *Ambio*, 20 (8), 366-371.
- Scoones, I., 1992. Coping with drought: responses of herders and livestock in contrasting Savanna environments in southern Zimbabwe. *Human Ecology*, 20 (3), 293-314.
- Scoones, I., 1994. *Living with uncertainty: new directions in pastoral development in Africa*. Intermediate Technology Publications, London.
- Scoones, I., 1995. Exploiting heterogeneity: habitat use by cattle in dryland Zimbabwe. *Journal of Arid Environments*, 29 (2), 221-237.

- Scott, C.B., Provenza, F.D. and Banner, R.E., 1995. Dietary habits and social interactions affect choice of feeding location by sheep. *Applied Animal Behaviour Science*, 45 (3/4), 225-237.
- Scott, L.L. and Provenza, F.D., 1998. Variety of foods and flavors affects selection of foraging location by sheep. *Applied Animal Behaviour Science*, 61 (2), 113-122.
- Scott, L.L. and Provenza, F.D., 1999. Variation in food selection among lambs: effects of basal diet and foods offered in a meal. *Journal of Animal Science*, 77 (9), 2391-2397.
- Scott, L.L. and Provenza, F.D., 2000. Lambs fed protein or energy imbalanced diets forage in locations and on foods that rectify imbalances. *Applied Animal Behaviour Science*, 68 (4), 293-305.
- Sellers, P.J., 1985. Canopy reflectance, photosynthesis and transpiration. *International Journal of Remote Sensing*, 6 (8), 1335-1372.
- Sellers, P.J., Los, S.O., Tucker, C.J., et al., 1996. A revised land surface parameterization (SiB2) for atmospheric GCMs. Part II: The generation of global fields of terrestrial biophysical parameters from satellite data. *Journal of Climate*, 9 (4), 706-737.
- Sellers, P.J., Tucker, C.J., Collatz, G.J., et al., 1994. A global 1° by 1° NDVI data set for climate studies. Part 2: The generation of global fields of terrestrial biophysical parameters from the NDVI. *International Journal of Remote Sensing*, 15 (17), 3519-3545.
- Senft, R.L., 1989. Hierarchical foraging models: effects of stocking and landscape composition on simulated resource use by cattle. *Ecological Modelling*, 46 (3/4), 283-303.
- Senft, R.L., Coughenour, M.B., Bailey, D.W., et al., 1987a. Large herbivore foraging and ecological hierarchies. *BioScience*, 37 (11), 789-799.
- Senft, R.L., Rittenhouse, L.R. and Woodmansee, D.R.G., 1985a. Factors influencing patterns of cattle grazing behavior on shortgrass steppe. *Journal of Range Management*, 38 (1), 82-87.
- Senft, R.L., Rittenhouse, L.R. and Woodmansee, R.G., 1985b. Factors influencing selection of resting sites by cattle on shortgrass steppe. *Journal of Range Management*, 38 (4), 295-299.
- Senft, R.L., Stillwell, M.A. and Rittenhouse, L.R., 1987b. Nitrogen and energy budgets of free-roaming cattle. *Journal of Range Management*, 40, 421.
- Shipley, L.A., 1999. Grazers and browsers: how digestive morphology affects diet selection. In: Launchbaugh, K.L., Sanders, K.D. and Mosley, J.C. eds. *Grazing behavior of livestock and wildlife*. University of Idaho, Moscow, 20-27. Idaho Forest, Wildlife and Range Experimental Station Bulletin no. 70.
- Shipley, L.A., Gross, J.E., Spalinger, D.E., et al., 1994. The scaling of intake rate in mammalian herbivores. *American Naturalist*, 143 (6), 1055-1082.
- Shipley, L.A. and Spalinger, D.E., 1992. Mechanics of browsing in dense food patches: effects of plant and animal morphology on intake rate. *Canadian Journal of Zoology*, 70, 1743-1752.
- Shipley, L.A., Spalinger, D.E., Gross, J.E., et al., 1996. The dynamics and scaling of foraging velocity and encounter rate in mammalian herbivores. *Functional Ecology*, 10 (2), 234-244.
- Shiyomi, M., 1987. Characteristics of herbage biomass estimated by the quadrat method. *Journal of Japanese Society of Grassland Science*, 32 (4), 307-312.
- Shiyomi, M., Akiyama, T. and Takahashi, S., 1983. A spatial pattern model of plant biomass in grazing pasture. *Journal of Japanese Society of Grassland Science*, 28 (4), 373-382.
- Shiyomi, M., Akiyama, T. and Takahashi, S., 1984. A spatial pattern model of plant biomass in grazing pasture. II. *Journal of Japanese Society of Grassland Science*, 30 (1), 40-48.
- Shiyomi, M., Koyama, N. and Tsuiki, M., 1991. Method for predicting available herbage biomass on grazed pasture. *Journal of Japanese Society of Grassland Science*, 36 (4), 327-337.
- Shiyomi, M., Okada, M., Takahashi, S., et al., 1998. Spatial pattern changes in aboveground plant biomass in a grazing pasture. *Ecological Research*, 13 (3), 313-322.
- Short, J., 1985. The functional response of kangaroos, sheep and rabbits in an arid grazing system. *The Journal of Applied Ecology*, 22 (2), 435-447.
- Sibly, R.M. and Smith, R.H., 1985. *Behavioural ecology: ecological consequences of adaptive behaviour: the 25th symposium of the British Ecological Society, Reading 1984*. Blackwell, Oxford.
- Silver, H., Colovos, N.F., Holter, J.B., et al., 1969. Fasting metabolism of white-tailed deer. *The Journal of Wildlife Management*, 33 (3), 490-498.
- Simpson, S.J. and Raubenheimer, D., 1993. A multi-level analysis of feeding behaviour: the geometry of nutritional decisions. *Philosophical Transactions: Biological Sciences*, 342 (1302), 381-402.
- Simpson, S.J. and Raubenheimer, D., 1999. Assuaging nutritional complexity: a geometrical approach. *Proceedings of the Nutrition Society*, 58 (4), 779-789.
- Simpson, S.J., Sibly, R.M., Lee, K.P., et al., 2004. Optimal foraging when regulating intake of multiple nutrients. *Animal Behaviour*, 68 (6), 1299-1311.
- Sinclair, A.R., Mduma, S. and Brashares, J.S., 2003. Patterns of predation in a diverse predator-prey system. *Nature*, 425 (6955), 288-290.

- Sinclair, A.R.E., 1977. *African buffalo: a study of resource limitation of populations*. University of Chicago Press, Chicago.
- Sinclair, A.R.E., 1985. Does interspecific competition or predation shape the African ungulate community? *Journal of Animal Ecology*, 54 (3), 899-918.
- Sinclair, A.R.E. and Arcese, P., 1995a. *Serengeti 2: dynamics, management, and conservation of an ecosystem [based on papers at a workshop held in December 1991 at the Serengeti Research Institute, Tanzania]*. University of Chicago Press, Chicago.
- Sinclair, A.R.E. and Arcese, P., 1995b. Population consequences of predation-sensitive foraging: the Serengeti wildebeest. *Ecology*, 76 (3), 882-891.
- Sinclair, A.R.E. and Norton-Griffiths, M., 1982. Does competition or facilitation regulate migrant ungulate populations in the Serengeti? A test of hypotheses. *Oecologia*, 53 (3), 364-369.
- Sinclair, A.R.E. and Smith, N.M., 1984. Do plant secondary compounds determine feeding preferences of snowshoe hares? *Oecologia*, 61 (3), 403-410.
- Skidmore, A.K., 1989. An expert system classifies eucalypt forest types using Thematic Mapper data and a digital terrain model. *PEandRS, Photogrammetric Engineering and Remote Sensing*, 55 (10), 1449-1464.
- Skidmore, A.K., Bijker, W., Schmidt, K., et al., 1997. Use of remote sensing and GIS for sustainable land management. *International Journal of Applied Earth Observation and Geoinformation* 3(4), 302-315.
- Skidmore, A.K., Schmidt, K.S., Kloosterman, H., et al., 2001. *Hyperspectral imagery for coastal wetland vegetation mapping*. Netherlands Remote Sensing Board (BCRS) Programme Bureau Rijkswaterstaat Survey Department, Delft.
- Skidmore, A.K., Watford, F., Luckananurug, P., et al., 1996. An operational GIS expert system for mapping forest soils. *Photogrammetric Engineering and Remote Sensing*, 62 (5), 501-511.
- Smith, N.S., 1970. Appraisal of condition estimation models for East African ungulates. *East African Wildlife Journal* 8, 123-129.
- Solanki, G.S., 1994. Feeding habits and grazing behavior of goats in a semi-arid region of India. *Small Ruminant Research*, 14 (1), 39-43.
- Spalinger, D.E., Hanley, T.A. and Robbins, C.T., 1988. Analysis of the functional response in foraging in the Sitka black-tailed deer. *Ecology*, 69 (4), 1166-1175.
- Spalinger, D.E. and Hobbs, N.T., 1992. Mechanisms of foraging in mammalian herbivores: new models of functional response. *The American Naturalist*, 140 (2), 325-348.
- Sperling, L. and Galaty, J.G., 1990. Cattle, culture, and economy: dynamics in East African pastoralism. In: Galaty, J.G. and Johnson, D.L. eds. *World of pastoralism: herding systems in comparative perspective*. Guildford Press, New York, 69-98.
- Stanley Price, M.R., 1974. *The feeding ecology of Coke's hartebeest, Alcelaphus buselaphus cokei Günther, in Kenya*. University of Oxford, Oxford. DPhil thesis
- Stein, A., Van Lieshout, M.N.M. and Boontink, H.W.G., 2001. Spatial interaction of methylene blue stained soil pores. *Geoderma*, 102 (1/2), 101-121.
- Stenning, D.J., 1957. Transhumance, migratory drift, migration: patterns of pastoral Fulani nomadism. *The Journal of the Royal Anthropological Institute of Great Britain and Ireland*, 87 (1), 57-73.
- Stenning, D.J., 1959. *Savannah nomads: a study of the Wodaabe pastoral Fulani of Western Bornu province Northern region, Nigeria*. Oxford University Press, Oxford.
- Stephens, D.W. and Krebs, J.R., 1986. *Foraging theory*. Princeton University Press, Princeton.
- Stobbs, T.H., 1973. The effect of plant structure on the intake of tropical pastures. I. Variation in the bite size of grazing cattle. *Australian Journal of Agricultural Research*, 24 (6), 809-819.
- Stone, L., Dayan, T. and Simberloff, D., 2000. On desert rodents, favored states, and unresolved issues: scaling up and down regional assemblages and local communities. *The American Naturalist*, 156 (3), 322-328.
- Stoyan, D., Kendall, W.S. and Mecke, J., 1995. *Stochastic geometry and its applications*. 2nd edn. Wiley, Chichester.
- Stricklin, W.R., 2001. The evolution and domestication of social behaviour. In: Keeling, L.J. and Gonyou, H.W. eds. *Social behaviour in farm animals*. Cabi, Wallingford, 83-112. [http://www.cabi.org/bk_BookDisplay.asp?PID=1487]
- Sullivan, S. and Rohde, R., 2002. On non-equilibrium in arid and semi-arid grazing systems. *Journal of Biogeography*, 29 (12), 1595-1618.
- Suttie, J.M., 2000. *Country pasture/forage resource profiles: Mongolia*. Food and Agriculture Organization of the United Nations, Rome. [<http://www.fao.org/ag/AGP/AGPC/doc/Counprof/mongol1.htm>]
- Swain, T., 1979. Tanins and lignins. In: Rosenthal, G.A. and Janzen, D.H. eds. *Herbivores: their interaction with secondary plant metabolites*. Academic Press, New York, 657-683.

- Swift, J., 1977. Sahelian pastoralists: underdevelopment, desertification, and famine. *Annual Review of Anthropology*, 6 (1), 457-478.
- Swift, J., 1986. The economics of production and exchange in West African pastoral societies. In: Adamu, M. and Kirk-Greene, A.H.M. eds. *The pastoralists of the West African savanna*. Manchester University Press, Manchester, 175-189.
- Sylla, D., 1994. Pastoral organizations for uncertain environments. In: Scoones, I. ed. *Living with uncertainty: new directions in pastoral development in Africa*. Intermediate Technology Publications, London, 134-152.
- Taylor, C.R., 1969. The eland and the oryx. *Scientific American*, 220 (1), 88-95.
- Taylor, R.J., 1984. Foraging in the eastern grey kangaroo and the wallaroo. *The Journal of Animal Ecology*, 53 (1), 65-74.
- Thornton, P.K., Galvin, K.A. and Boone, R.B., 2003. An agro-pastoral household model for the rangelands of East Africa. *Agricultural Systems*, 76 (2), 601-622.
- Tieleman, B.I. and Williams, J.B., 2000. The adjustment of avian metabolic rates and water fluxes to desert environments. *Physiological and Biochemical Zoology*, 73 (4), 461-479.
- Tilman, D., 1982. *Resource competition and community structure*. Princeton University Press, Princeton.
- Tilman, D., 1988. *Plant strategies and the dynamics and structure of plant communities*. Princeton University Press, Princeton.
- Tilman, D. and Kareiva, P., 1997. *Spatial ecology: the role of space in population dynamics and interspecific interactions*. Princeton University Press, Princeton.
- Trudell, J. and White, R.G., 1981. The effect of forage structure and availability on food intake, biting rate, bite size and daily eating time of reindeer. *Journal of Applied Ecology*, 18 (1), 63-81.
- Tucker, C.J., 1977. Asymptotic nature of grass canopy spectral reflectance. *Applied Optics*, 16 (5), 1151-1156.
- Tucker, C.J., Vanpraet, C.L., Sharman, M.J., et al., 1985. Satellite remote sensing of total herbaceous biomass production in the Senegalese Sahel: 1980-1984. *Remote Sensing of Environment*, 17, 233-249.
- Turchin, P., 2003. *Complex population dynamics: a theoretical/empirical synthesis*. Princeton University Press, Princeton.
- Turner, M.D., 1998. Long-term effects of daily grazing orbits on nutrient availability in Sahelian West Africa: I. Gradients in the chemical composition of rangeland soils and vegetation. *Journal of Biogeography*, 25 (4), 669-682.
- Turner, M.D., 1999a. Merging local and regional analyses of land-use change: the case of livestock in the Sahel. *Annals of the Association of American Geographers*, 89 (2), 191-219.
- Turner, M.D., 1999b. The role of social networks, indefinite boundaries and political bargaining in maintaining the ecological and economic resilience of the transhumance systems of Sudano-Saharan West Africa. In: Niamir-Fuller, M. ed. *Managing mobility in African rangelands: the legitimization of transhumance*. Intermediate Technology Publications, London, 97-123.
- Turner, M.D., 2003. Multiple holders of multiple stakes: the multilayered politics of agropastoral resource management in semi-arid Africa. In: Allsopp N, P.A., Milton SJ, Kirkman KP, Kerley GIH, Hurt CR, Brown CJ ed. *Proceedings of the VIIIth International Rangeland Congress, July 26-August 1, 2003, Durban, South Africa*. 1715-1725.
- Turner, M.G., Wu, Y., Romme, W.H., et al., 1993. A landscape simulation model of winter foraging by large ungulates. *Ecological Modelling*, 69 (3/4), 163-184.
- Tyler, N.J.C., 1987. Body composition and energy balance of pregnant and non-pregnant Svalbard reindeer during winter. *Symposia of the Zoological Society of London*, 57, 203-229.
- Underwood, R., 1983. The feeding behaviour of grazing African ungulates. *Behaviour*, 84 (3/4), 195-243.
- UNEP, 1992. *World atlas of desertification*. Arnold, London.
- Ungar, E.D., 1996. Ingestive behaviour. In: Hodgson, J. and Illius, A.W. eds. *The ecology and management of grazing systems*. CAB international, Wallingford, 185-218.
- Ungar, E.D., Genizi, A. and Demment, M.W., 1991. Bite dimensions and herbage intake by cattle grazing short hand-constructed swards. *Agronomy Journal*, 83 (6), 973-978.
- Valentine, K.A., 1947. Distance from water as a factor in grazing capacity of rangeland. *Journal of Forestry*, 45, 749-754.
- Vallentine, J.F., 2001. *Grazing management*. 2nd edn. Academic Press, San Diego.
- Van de Koppel, J., 1997. *Trophic interactions in gradients of primary productivity*. University of Groningen, Groningen. PhD thesis
- Van de Koppel, J., Huisman, J., Van der Wal, R., et al., 1996. Patterns of herbivory along a productivity gradient: an empirical and theoretical investigation. *Ecology*, 77 (3), 736-745.

- Van de Koppel, J., Rietkerk, M., Van Langevelde, F., et al., 2002. Spatial heterogeneity and irreversible vegetation change in semiarid grazing systems. *American Naturalist*, 159 (2), 209-218.
- Van de Vijver, C., Poot, P. and Prins, H.H.T., 1999. Causes of increased nutrient concentrations in post-fire regrowth in an East African savanna. *Plant and Soil*, 214 (1), 173-185.
- Van de Vijver, C.A.D.M., 1999. *Fire and life in Tarangire: effects of burning and herbivory on an East African Savanna system*. Wageningen University, Wageningen. PhD thesis
- Van der Jeugd, H.P. and Prins, H.H.T., 2000. Movements and group structure of giraffe (*Giraffa camelopardalis*) in Lake Manyara National Park, Tanzania. *Journal of Zoology*, 251 (01), 15-21.
- Van der Wal, R., Madan, N., Van Lieshout, S., et al., 2000. Trading forage quality for quantity? Plant phenology and patch choice by Svalbard reindeer. *Oecologia*, 123 (1), 108-115.
- Van Hoven, W. and Frustenburg, D., 1992. The use of purified condensed tannin as a reference in determining its influence on rumen fermentation. *Comparative Biochemistry and Physiology A, Comparative Physiology*, 101 (2), 381-385.
- Van Keulen, H. and Breman, H., 1990. Agricultural development in the West African Sahelian region: a cure against land hunger? *Agriculture, Ecosystems and Environment*, 32 (3/4), 177-197.
- Van Langevelde, F., Van de Vijver, C.A.D.M., Lalit, K., et al., 2003. Effects of fire and herbivory on the stability of savanna ecosystems. *Ecology*, 84 (2), 337-350.
- Van Lieshout, M.N.M. and Baddeley, A.J., 1999. Indices of dependence between types in multivariate point patterns. *Scandinavian Journal of Statistics*, 26 (4), 511-532.
- Van Soest, P.J., 1982. *Nutritional ecology of the ruminant: ruminant metabolism, nutritional strategies, the cellulolytic fermentation and the chemistry of forages and plant fibers*. O & B Books, Corvallis.
- Van Soest, P.J., 1994. *Nutritional ecology of the ruminant*. 2nd edn. Comstock, Ithaca.
- Van Wieren, S.E., 1992. Factors limiting food intake in ruminants and non-ruminants in the temperate zone. In: Spitz, F., Janeau, G., Gonzalez, G., et al. eds. *Ongules - ungulates: proceedings of the international symposium "Ongules - ungulates", Toulouse, France, September 2-6, 1991*. S.F.E.P.M., Paris, 139-145.
- Van Wieren, S.E., 1996a. *Digestive strategies in ruminants and nonruminants*. Wageningen University, Wageningen. PhD thesis
- Van Wieren, S.E., 1996b. Do large herbivores select a diet that maximizes short-term energy intake rate? *Forest Ecology and Management*, 88 (1/2), 149-156.
- Vavra, M. and Ganskopp, D., 1998. Grazing behavior in ungulates: current concepts and future challenges. *Annals of Arid Zones*, 37 (3), 319-335.
- Verheyden-Tixier, H. and Duncan, P., 2000. Selection for small amounts of hydrolysable tannins by a concentrate-selecting mammalian herbivore. *Journal of Chemical Ecology*, 26 (2), 351-358.
- Verweij, R.J.T., Verrelst, J., Loth, P.E., et al., 2006. Grazing lawns contribute to the subsistence of mesoherbivores on dystrophic savannas. *Oikos*, 114 (1), 108-116.
- Vesey-FitzGerald, D.F., 1974. Utilisation of grazing resources by buffalo in the Arusha National Park, Tanzania. *East African Wildlife Journal* 12, 107-134.
- Villalba, J.J. and Provenza, F.D., 1996. Preference for flavored wheat straw by lambs conditioned with intraruminal administrations of sodium propionate. *Journal of Animal Science*, 74 (10), 2362-2368.
- Villalba, J.J. and Provenza, F.D., 1997a. Preference for flavored wheat straw by lambs conditioned with intraruminal infusions of acetate and propionate. *Journal of Animal Science*, 75 (11), 2905-2914.
- Villalba, J.J. and Provenza, F.D., 1997b. Preference for flavoured foods by lambs conditioned with intraruminal administration of nitrogen. *British Journal of Nutrition*, 78 (4), 545-561.
- Villalba, J.J. and Provenza, F.D., 1997c. Preference for wheat straw by lambs conditioned with intraruminal infusions of starch. *British Journal of Nutrition*, 77 (2), 287-297.
- Villalba, J.J. and Provenza, F.D., 1999. Nutrient-specific preferences by lambs conditioned with intraruminal infusions of starch, casein, and water. *Journal of Animal Science*, 77 (2), 378-387.
- Villalba, J.J. and Provenza, F.D., 2002. Polyethylene glycol influences selection of foraging location by sheep consuming quebracho tannin I. *Journal of Animal Science*, 80 (7), 1846-1851.
- VITO, 2003. *VEGETATION*. VITO. Available: [<http://www.vgt.vito.be/indexstart.htm>].
- Voeten, M.M., 1999. *Living with wildlife: coexistence of wildlife and livestock in an East African Savanna system*. Wageningen University, Wageningen. PhD thesis
- Voeten, M.M. and Prins, H.H.T., 1999. Resource partitioning between sympatric wild and domestic herbivores in the Tarangire region of Tanzania. *Oecologia*, 120 (2), 287-294.
- Vogelmann, J. and Moss, D., 1993. Spectral reflectance measurements in the genus *Sphagnum*. *Remote Sensing of Environment*, 45 (3), 273-279.
- Vos, J. and Bom, M., 1993. Hand-held chlorophyll meter: a promising tool to assess the nitrogen status of potato foliage. *Potato Research*, 36 (4), 301-308.

- Vucetich, J.A., Peterson, R.O. and Schaefer, C.L., 2002. The effect of prey and predator densities on wolf predation. *Ecology*, 83 (11), 3003-3013.
- Wackernagel, H., 1995. *Multivariate geostatistics: an introduction with applications*. Springer, Berlin.
- Wagenaar, K.T., Diallo, A. and Sayers, A.R., 1986. *Productivity of transhumant Fulani cattle in the inner Niger delta of Mali*. ILCA, Addis Ababa.
- Walker, B.H., Emslie, R.H., Owen-Smith, R.N., et al., 1987. To cull or not to cull: lessons from a Southern African drought. *The Journal of Applied Ecology*, 24 (2), 381-401.
- Walker, J., Jupp, D.L.B., Penridge, L.K., et al., 1986. Interpretation of vegetation structure in Landsat MSS imagery: a case study in disturbed semi-arid eucalypt woodlands. Part 1. Field data analysis. *Journal of Environmental Management*, 23 (1), 19-33.
- Walker, P.A., 1990. Modelling wildlife distributions using a geographic information system: kangaroos in relation to climate. *Journal of Biogeography*, 17 (3), 279-289.
- Wallace, J.S., Brouwer, J. and Allen, S. J., 1994. *Hapex-Sahel southern super-site report: an overview of the Hapex-Sahel southern super-site and the experimental programme during the intensive observation period in 1992*. Institute of Hydrology, Wallingford.
- Wallis de Vries, M.F., 1996. Effects of resource distribution patterns on ungulate foraging behaviour: a modelling approach. *Forest Ecology and Management*, 88 (1/2), 167-177.
- Wallis de Vries, M.F. and Daleboudt, C., 1994. Foraging strategy of cattle in patchy grassland. *Oecologia*, 100 (1/2), 98-106.
- Wallis de Vries, M.F., Laca, E.A. and Demment, M.W., 1999. The importance of scale of patchiness for selectivity in grazing herbivores. *Oecologia*, 121 (3), 355-363.
- Wallis de Vries, M.F. and Schippers, P., 1994. Foraging in a landscape mosaic: selection for energy and minerals in free-ranging cattle. *Oecologia*, 100 (1/2), 107-117.
- Wallis, I.R., Green, B. and Newgrain, K., 1997. Seasonal field energetics and water fluxes of the long-nosed potoroo (*Potorous tridactylus*) in southern Victoria. *Australian Journal of Zoology*, 45 (1), 1-11.
- Walter, H., 1971. *Ecology of tropical and subtropical vegetation*. Oliver and Boyd, Edinburgh.
- Wang, J. and Provenza, F.D., 1996. Food deprivation affects preference of sheep for foods varying in nutrients and a toxin. *Journal of Chemical Ecology*, 22 (11), 2011-2021.
- Ward, D., 1992. The role of satisficing in foraging theory. *Oikos*, 63 (2), 312-317.
- Ward, D. and Saltz, D., 1994. Foraging at different spatial scales: dorcas gazelles foraging for lilies in the Negev desert. *Ecology*, 75 (1), 48-58.
- Waser, P., 1975. Diurnal and nocturnal strategies of the bushbuck. *East African Wildlife Journal*, 13, 49-64.
- Weiher, E. and Keddy, P.A., 1999. Relative abundance and evenness patterns along diversity and biomass gradients. *Oikos*, 87 (2), 355-361.
- Weiner, J., 1977. Energy metabolism of the roe deer. *Acta Theriologica*, 22 (1/8), 3-24.
- West, G.B., Savage, V.M., Gillooly, J., et al., 2003. Why does metabolic rate scale with body size? *Nature*, 421 (6924), 713.
- West, G.B., Woodruff, W.H. and Brown, J.H., 2002. Allometric scaling of metabolic rate from molecules and mitochondria to cells and mammals. *PNAS*, 99 (Suppl. 1), 2473-2478.
- Westoby, M., 1978. What are the biological bases of varied diets? *The American Naturalist*, 112 (985), 627-631.
- Westoby, M., 1979. Elements of a theory of vegetation dynamics in arid rangelands. *Israel Journal of Botany*, 28 (3/4), 169-194.
- Westoby, M., Walker, B. and Noy-Meir, I., 1989. Opportunistic management for rangelands not at equilibrium. *Journal of Range Management*, 42 (4), 266-274.
- Wickstrom, M.L., Robbins, C.T., Hanley, T.A., et al., 1984. Food intake and foraging energetics of elk and mule deer. *Journal of Wildlife Management*, 48 (4), 1285-1301.
- Wiens, J.A., 1992. *The ecology of bird communities*. Repr. edn. Cambridge University Press, Cambridge.
- Williams, J.B., Anderson, M.D. and Richardson, P.R.K., 1997. Seasonal differences in field metabolism, water requirements, and foraging behavior of free-living aardwolves. *Ecology*, 78 (8), 2588-2602.
- Williams, R.E., 1954. Modern methods of getting uniform use of ranges. *Journal of Range Management*, 7 (2), 77-81.
- Wilmshurst, J.F. and Fryxell, J.M., 1995. Patch selection by red deer in relation to energy and protein intake: a re-evaluation of Langvatn and Hanley's (1993) results. *Oecologia*, 104 (3), 297-300.
- Wilmshurst, J.F., Fryxell, J.M. and Bergman, C.M., 2000. The allometry of patch selection in ruminants. *Proceedings of the Royal Society of London Series B, Biological Sciences*, 267 (1441), 345-349.
- Wilmshurst, J.F., Fryxell, J.M. and Colucci, P.E., 1999a. What constrains daily intake in Thomson's gazelles? *Ecology*, 80 (7), 2338-2347.

- Wilmshurst, J.F., Fryxell, J.M., Farm, B.P., et al., 1999b. Spatial distribution of Serengeti wildebeest in relation to resources. *Canadian Journal of Zoology*, 77 (8), 1223-1232.
- Wilmshurst, J.F., Fryxell, J.M. and Hudson, R.J., 1995. Forage quality and patch choice by wapiti (*Cervus elaphus*). *Behavioral Ecology*, 6 (2), 209-217.
- Wilson, E.O. and Bossert, W.H., 1971. *A primer of population biology*. Sinauer, Sunderland.
- Wilson, R.T., 1980. Fuelwood consumption in a central Malian town and its effects on browse availability. In: Le Houerou, H.N. ed. *Browse in Africa: the current state of knowledge*. International Livestock Centre for Africa, Addis Ababa, 473-476.
- Wilson, S.L. and Kerley, G.I.H., 2003. Bite diameter selection by thicket browsers: the effect of body size and plant morphology on forage intake and quality. *Forest Ecology and Management*, 181 (1/2), 51-65.
- Winston, W.L., 1994. *Operations research: applications and algorithms*. Duxbury, Belmont.
- Wint, W., 1998. Rapid resource assessment and environmental monitoring using low level aerial surveys. In: Squires, V.R. and Sidahmed, A.E. eds. *Drylands: sustainable use of rangelands into the twenty-first century*. IFAD, Rome, 277-301.
- Winterbach, H.E.K. and Bothma, J.D.P., 1998. Activity patterns of the Cape buffalo *Syncerus caffer caffer* in the Willem Pretorius Game Reserve, Free State. *South African Journal of Wildlife Research*, 28 (3), 73-81.
- Wissinger, S.A., Sparks, G.B., Rouse, G.L., et al., 1996. Intraguild predation and cannibalism among larvae of detritivorous caddisflies in subalpine wetlands. *Ecology*, 77 (8), 2421-2430.
- Woodward, F.I., 1987. *Climate and plant distribution*. Cambridge University Press, Cambridge.
- Worden, J., Reid, R. and Gichohi, H., 2003. *Land-use impacts on large wildlife and livestock in the swamps of the greater Amboseli ecosystem, Kajiado District, Kenya*. International Livestock Research Institute Nairobi. Land Use Change Impacts and Dynamics (LUCID) Working Paper no. 27.
- Wyant, J.G. and Ellis, J.E., 1990. Compositional patterns of riparian woodlands in the Rift Valley of northern Kenya. *Vegetatio*, 89 (1), 23-37.
- Wynne-Edwards, V.C., 1962. *Animal dispersion in relation to social behaviour*. Oliver and Boyd, Edinburgh.
- Ydenberg, R.C., Prins, H.H.T. and Van Dijk, J., 1983. Postroost gatherings of wintering Barnacle Geese: information centres. *Ardea*, 73, 125-132.
- Yearsley, J., Hastings, I.M., Gordon, I.J., et al., 2002. A lifetime perspective on foraging and mortality. *Journal of Theoretical Biology*, 215 (4), 385-397.
- Zervas, G., 1998. Quantifying and optimizing grazing regimes in Greek mountain systems. *Journal of Applied Ecology*, 35 (6), 983-986.
- Zonneveld, I.S., 1988. Survey approaches. In: Küchler, A.W. and Zonneveld, I.S. eds. *Vegetation mapping*. Kluwer, Dordrecht.