

Effect of sward age on nitrate accumulation in ryegrass

A. Darwinkel

Research Station for Arable Farming, Edelhertweg 1, Lelystad, the Netherlands

Accepted: 1 September 1976

Key words: nitrate content, nitrate accumulation, nitrate conversion, sward age, *Lolium perenne*, *Lolium multiflorum*

Summary

Two indoor trials were done with Italian ryegrass (*Lolium multiflorum*) and perennial ryegrass (*Lolium perenne*) to obtain more information about the nitrate-accumulating process of a new-sown grass sward.

With an abundant NO_3 supply for Italian ryegrass, NO_3 accumulation in herbage clearly decreased during five successive cuts after sowing, because of an increasing NO_3 conversion. In the same way the NO_3 content in dry matter in herbage of perennial ryegrass was considerably higher in the first cut after sowing than in old plants from a 5-year-old pasture.

After sowing, a considerable part of the assimilate produced was used for stubble and root growth. With older swards these assimilates were increasingly retained in herbage, which is rich in organic N and to a lesser extent in stubble and roots, which are poor in organic N. Because of this, NO_3 conversion is restricted in a new-sown grass sward and reaches to a maximum in old sward of a permanent pasture, which almost exclusively produces herbage. Consequently nitrate is more apt to accumulate in new-sown grass than in old pasture.

Introduction

As an important nutrient element, nitrogen has received much attention during research on how to increase dry matter yields. Maximum yields are only achieved with a sufficient supply of nitrogen. In grasses, this occurs in the presence of about 0.14 % $\text{NO}_3\text{-N}$ in dry matter (van Burg, 1965), but large amounts of NO_3 in the plant are unhealthy for cattle.

Large contents of NO_3 occur in various crops, as in oats (Gilbert et al., 1946) and more recently in stubble turnips (te Velde, 1967). Many articles describe factors that influence the NO_3 content of the plant. Light intensity, drought, N fertilizer and plant species are often mentioned (Deinum, 1966).

In herbage from permanent pasture, high contents of NO_3 hardly occur, not even

EFFECT OF SWARD AGE ON NITRATE ACCUMULATION IN RYEGRASS

after a large nitrogen dressing (de Groot et al., 1972). However, high NO_3 contents have been found in herbage from newly sown grasses (Dapper & Keuning, 1972; Thomas & Willemsen, 1971). These high contents were attributed to heavy dressing with fertilizer and farmyard manure, but with similar fertilization the NO_3 content of newly sown swards was considerably higher than that of older swards. So the question arises whether the newly sown swards themselves are a factor in increasing NO_3 content.

This paper summarizes the results of two indoor trials, in which Kjeldahl N and nitrate N were estimated in (i) five successive cuts of Italian ryegrass and (ii) new-sown and old perennial ryegrass. So the possible specific behaviour on nitrate accumulation of newly sown swards can be studied.

Materials and methods

The pot trials were in the phytotron of the Department of Field Crops and Grassland Husbandry, Agricultural University at Wageningen.

Trial 1. Dry matter production and nitrate accumulation in successive cuts of Italian ryegrass (Lolium multiflorum)

0.6 g seed of Italian ryegrass cv. Tetila was sown on a layer of fine gravel and placed on pots, containing 5 litres of a nutrient solution. The pots were placed in a growth cabinet at 16/12 °C (day/night). Daylength was 15 h and daily irradiance was about $800 \text{ J cm}^{-2} \text{ d}^{-1}$ (400 - 700 nm).

During the trial, the grass was cut five times at intervals of 3 weeks. After each growth period, the herbage was harvested and the nutrient solution renewed. The two N treatments were 400 and 800 mg N per pot after each cut. Each treatment included 5 pots per cut.

Trial 2. Effect of plant age on dry matter production and nitrate accumulation in perennial ryegrass (Lolium perenne)

On 30 September 1970, 5-litre plastic pots were filled with sandy soil and were kept in a temperature-controlled greenhouse with 4 replicates at 20/15 °C (day/night). Plants of a 5-year-old pasture of perennial ryegrass sown as a mixture of cv. Barenza and cv. Pelo, 1 : 1, were planted in half the pots on 6 October. On 9 November the other pots were sown with 0.9 g of the same mixture of grass seed. At the beginning of the trial on 23 November the old plants were cut for a second time, when the sown grass showed a good initial development. Afterwards 400 or 1000 mg N per pot were added. On 21 December, after 4 weeks, the plants were harvested.

Daylength was 14 h and average irradiance, including that of HPLR lamps, was about $950 \text{ J cm}^{-2} \text{ d}^{-1}$ (400 - 700 nm).

Analysis

To obtain more information on the nitrate-accumulating process, harvested plants were separated into herbage, stubble and roots, since these tissues widely differ in nitrate metabolism (Darwinkel, 1975). After drying at 80 °C for at least 20 h, the

dried material was ground through a 1-mm sieve and stored in air-tight plastic bags for analysis.

For total nitrogen (including nitrate), the Kjeldahl method as modified by Deys (1961) was used. The content of NO_3 was estimated potentiometrically with a nitrate-specific electrode.

Results

Dry matter production

The production and distribution of dry matter was recorded by weighing herbage, stubble and roots. Table 1 presents the results of both trials and Fig. 1 schematizes data from the treatment with heavy N dressing.

In Trial 1, dry matter yield increased with each successive cut, and reached a maximum at the 4th cut. However in Trial 2, dry matter production of the old plants was considerably lower than of young plants, because of a small initial rate of regrowth of the old plants after cutting on 23 November.

Table 1 shows the total dry matter of each cut in Trial 1 was less than in Trial 2, mainly because of the shorter period of growth. In both trials dry matter increased with N supply.

Table 1 also shows that herbage production was much greater than production of stubble + roots in each growth period. In Trial 1, yield of herbage increased at each successive cut, but of stubble + roots was almost constant: an increasing proportion of production from sown grass was retained in herbage. At the 1st cut, more than 40 % of dry matter was used for growth of roots and stubble, but at the 5th cut this was less than 20 %.

In Trial 2, the differences in distribution of dry matter were even greater. The young plants laid down a considerable proportion of dry matter in stubble and roots,

Table 1. Increase of dry matter (g per pot) of grasses at 2 nitrate dressings. A. Trial 1: newly sown plants of Italian ryegrass during 5 successive cuts. B. Trial 2: newly sown and old plants of perennial ryegrass.

A. Trial 1	Total dry matter		Herbage		Stubble + roots	
	400 mg N	800 mg N	400 mg N	800 mg N	400 mg N	800 mg N
1st cut	11.00	10.77	6.14	6.43	4.86	4.31
2nd cut	11.79	13.42	6.85	10.10	3.14	3.32
3rd cut	12.66	14.98	9.82	12.62	2.84	2.36
4th cut	15.93	18.94	11.48	13.89	4.45	5.05
5th cut	16.25	18.73	12.81	15.96	3.44	2.77
B. Trial 2	Total dry matter		Herbage		Stubble + roots	
	400 mg N	1000 mg N	400 mg N	1000 mg N	400 mg N	1000 mg N
New-sown plants	21.04	22.12	12.88	14.53	8.16	7.59
Old plants	15.79	16.75	14.37	16.13	1.42	0.62

EFFECT OF SWARD AGE ON NITRATE ACCUMULATION IN RYEGRASS




Fig. 1. Increase of dry matter (g per pot) of grasses with high nitrate supply. Left: Trial 1 with Italian ryegrass during 5 successive cuts, 800 mg N per pot. Right: Trial 2 with new-sown and old perennial ryegrass, 1000 mg N per pot.

Table 2. Nitrate contents (g nitrate N/100 g dm) in herbage, stubble and roots of Italian ryegrass in 5 successive cuts (Trial 1) and of newly sown and old perennial ryegrass plants (Trial 2) at 2 N dressings.

<i>Trial 1</i>	Herbage		Stubble		Roots	
	400 mg N	800 mg N	400 mg N	800 mg N	400 mg N	800 mg N
1st cut	0.88	1.66	0.71	1.33	0.38	1.43
2nd cut	0.61	1.27	0.70	1.31	0.39	1.07
3rd cut	0.55	1.24	0.82	1.52	0.37	0.58
4th cut	0.31	0.93	0.44	1.41	0.11	0.38
5th cut	0.22	0.84	0.29	0.96	0.07	0.22
<i>Trial 2</i>	Herbage		Stubble		Roots	
	400 mg N	1000 mg N	400 mg N	1000 mg N	400 mg N	1000 mg N
New-sown plants	0.21	0.63	0.26	0.66	0.04	0.24
Old plants	0.15	0.33	0.14	0.29	0.06	0.12

whereas the old plants showed hardly any increase in stubble + roots. The increase in dry matter in old plants consisted mainly of herbage.

Nitrogen influenced distribution of dry matter in both trials. Herbage yield increased clearly with a higher nitrogen supply, but yield of stubble and roots was hardly affected.

Nitrate contents

Table 2 shows the NO_3 content of herbage, stubble and roots in both trials. For a good comparison of NO_3 contents, yield of dry matter must be known. In Trial 2, the NO_3 content was lower than in Trial 1, mainly because of the higher yield of dry matter. Clear differences in NO_3 content existed between herbage, stubble and roots. The lowest contents were always found in roots. At the first cut of Trial 1, herbage had a higher NO_3 content than stubble, but at successive cuts thereafter the reverse was found. In Trial 2, the differences in NO_3 content between herbage and stubble were small. As expected, NO_3 content in all plant parts increased with a higher N supply.

Plant age clearly influenced NO_3 content. In Trial 1, NO_3 content continued to decrease at each of the 5 successive cuts. This decrease was clear in herbage and roots, but less pronounced in stubble. The effect of plant age was also present in Trial 2, in which old plants had considerably less NO_3 than the young ones.

During the 5 consecutive growth periods of Trial 1, NO_3 content decreased by 80 % with the low dressing and by 50 % with the high dressing. Consequently the effect of N supply on NO_3 content was highest at the 5th harvest.

So the difference in NO_3 content between old and newly sown swards, as found in practice, occurred also in these trials, indicating that N supply is of little bearing of these differences.

Discussion

Accumulation of NO_3 in plants is the difference between uptake and conversion of NO_3 . Uptake depends closely upon the supply of nitrogen, whereas conversion depends upon production and distribution of dry matter (Darwinkel, 1975). Knowledge about dry matter production and about uptake and conversion of NO_3 is necessary to analyse the nitrate-accumulating process.

During initial growth of grasses after sowing, much assimilate is used for root and stubble development. In course of time, a dense sward and an extensive root system is produced. Gradually the weight of stubble and roots practically stabilizes. Then, stubble and roots will take little more assimilate than is necessary to compensate for ageing parts. Such an equilibrium is found in permanent pasture.

To achieve this equilibrium a gradual change takes place in the distribution of dry matter between herbage, stubble and roots. During ageing of the sward, an increasing proportion of the dry matter produced is retained in herbage. This change in dry matter distribution favours nitrate conversion as the content of organic N in herbage is considerably higher than in stubble and roots (Alberda, 1965; Darwinkel, 1975).

This equilibrium in distribution of dry matter was not yet reached in Trial 1 after 5 cuts (Table 1). During the first growth period after sowing, more than 40 % of dry matter was retained in roots and stubble. During successive periods of the growth, this share decreased gradually to less than 20 % at the 5th cut. Also in Trial 2, 40 % of the dry matter produced was in stubble and roots of the young plants, whereas weight of stubble and roots hardly changed in old plants (Table 3).

Yield increases of dry matter and nitrogen are presented in Table 3.

In Trial 1, the increments of dry matter in whole plant and in herbage were higher in each successive cut (Table 1). In Trial 2 increment of dry matter of old plants lagged behind that of young plants, because of the slow initial regrowth.

With the same abundant N supply, nitrogen yield at the 5th cut of Trial 1 was greater than at the 1st cut, possibly because of the more extensive root system. The conversion of NO_3 into organic N increased even more than N uptake, so that a

Table 3. Increment of dry matter (g) and of nitrogen (mg) per pot in Italian ryegrass at 1st and 5th cut (Trial 1) and in new-sown and old perennial ryegrass (Trial 2) with high N supply.

	Dry matter production			Total N	Nitrate N	Organic N
	total	herbage	stubble + roots			
<i>Trial 1</i>						
1st cut	10.74	6.43	4.31	568	166	402
5th cut	18.73	15.96	2.77	674	126	548
<i>Trial 2</i>						
New-sown plants	22.12	14.53	7.59	703	137	566
Old plants	16.75	16.13	0.62	629	64	565

Table 4. Amount (mg) and proportion of nitrate accumulated per pot in herbage and stubble + roots with high N supply in the two trials.

	Amount of nitrate (mg)		Proportion of nitrate (%)	
	herbage	stubble + roots	herbage	stubble + roots
<i>Trial 1</i>				
1st cut	107	59	65	35
5th cut	134	103	57	43
<i>Trial 2</i>				
New-sown plants	95	55	63	37
Old plants	51	43	54	46

smaller amount of NO_3 had been accumulated at the 5th cut. Moreover, this lower NO_3 accumulation was achieved at a higher yield of dry matter, so that the difference in NO_3 content between the 1st and 5th cut was very high (Table 2). In Trial 2 both groups of plants converted equal amounts of NO_3 into organic N, so the greater uptake by young plants led to greater accumulation.

Leaves have a considerably higher content of organic N than the other plant parts, so growth of leaves determines NO_3 conversion. In Trial 1 the greater yield of organic N at the 5th cut must mainly be attributed to a higher herbage yield. In Trial 2, the influence of herbage yield is even clearer. Although yield of dry matter in whole plants was less in old plants, yield of herbage was more than in young plants, and so both plants contained equal amounts of organic N.

So distribution of dry matter influences the conversion and accumulation of NO_3 . For morphologically different varieties of stubble turnip and for Italian and Westerwolths ryegrass, differences in NO_3 conversion depend upon differences in yield of leaves (Darwinkel, 1975). So differences in NO_3 content between species and varieties can be explained by differences in distribution of dry matter. If a considerable proportion of the dry matter produced accumulates in plant parts poor in organic N, such as roots and stems, NO_3 conversion is restricted. On the contrary, a large accumulation of dry matter in leaves stands for a good conversion. Perhaps low NO_3 contents in perennials and the high contents in annuals (Kretschmer, 1958) are connected with this difference in dry matter distribution.

Measurements of the activity of the enzyme nitrate reductase showed that NO_3 reduction occurs mainly in leaves (Wallace & Pate, 1967). In grasses, green stubble also contributes to some extent to NO_3 reduction (Darwinkel, 1975). Probably the great amount of stubble favoured NO_3 conversion during the later growth periods of Trial 1 and in the old plants of Trial 2.

High NO_3 contents rarely occur in permanent pasture, even with high N dressings (de Groot et al., 1972), possibly because dry matter is mainly produced as herbage, in which much NO_3 can be converted into organic compounds. Moreover, a considerable proportion of the NO_3 taken up can be accumulated in the large storage capacity of stubble and roots. Newly sown swards and annual grasses have only a limited capacity in stubble and roots and consequently a major part of the NO_3 is

EFFECT OF SWARD AGE ON NITRATE ACCUMULATION IN RYEGRASS

translocated to herbage. Table 4 presents distribution of NO_3 between herbage, stubble and roots. With ageing of the sward, an ever smaller proportion of the NO_3 is found in herbage. The high NO_3 content in stubble (Table 2) indicates that NO_3 accumulates there. With ageing, the amount of stubble increases and even more NO_3 can accumulate in it. Accordingly, less NO_3 is transported to herbage, whereas herbage yield increases with plant age (Fig. 1). This results in a sharp decrease in the NO_3 content.

Acknowledgments

The author is much indebted to Prof. Ir M. L. 't Hart and Dr Ir B. Deinum for valuable comments and to Mrs Evelyn Zantman for technical assistance.

References

- Alberda, Th., 1965. The influence of temperature, light intensity and nitrate concentration on dry matter production and chemical composition of *Lolium perenne* L. *Neth. J. agric. Sci.* 13: 335-360.
- Burg, P. F. J. van, 1965. De stikstofbemesting van grasland. 7. Het nitraatgehalte als indicator voor de stikstofvoeding. *Stikstof* 47/48: 461-469.
- Dapper, J. G. & J. A. Keuning, 1972. Hoge nitraatgehalten in het gras en in grasprodukten op het bedrijf Reynen te Gemert. *Stikstof* 71: 476-483.
- Darwinkel, A., 1975. Aspects of assimilation and accumulation of nitrate in some cultivated plants. *Agric. Res. Rep.* 843. Pudoc, Wageningen, 64 pp.
- Deinum, B., 1966. Climate, nitrogen and grass. Research into the influence of light intensity, temperature, water supply and nitrogen on the production and chemical composition of grass. *Meded. LandbHogeschool Wageningen* 66-11; 91 pp.
- Deys, W. B., 1961. De bepaling van totaal stikstof in gewasmonsters, inclusief nitraat-stikstof. *Jaarb. Inst. biol. scheik. Onderz. LandbGewassen*: 89-91.
- Gilbert, C. S., H. F. Eppson, W. B. Bradley & O. A. Beath, 1946. Nitrate accumulation in cultivated plants and weeds. *Bull. Univ. Wyo. agric. Exp. Stn* 277; 39 pp.
- Groot, Th. de, J. A. Keuning & L. Padmos, 1972. Hoge stikstofgiften op grasland en de produktie en gezondheid van melkvee. *Stikstof* 72: 495-532.
- Kretschmer, A. E., 1958. Nitrate accumulation in Everglades forages. *Agron. J.* 50: 314-316.
- Thomas, H. & W. Willemsen, 1971. Hoge nitraatgehalten in graslandprodukten. *Stikstof* 68: 313-319.
- Velde, H. A. te, 1967. Nitraatgehalten van stoppelknollen in 1966 (I). *Meded. Proefstn Akker- en Weideb.* 139; 46 pp.
- Wallace, W. & J. S. Pate, 1967. Nitrate assimilation in higher plants with special reference to the cocklebur (*Xanthium pennsylvanicum* Wallr.). *Ann. Bot.* 31: 213-228.