

- STOKES, W. E. and F. H. HULL : Peanut breeding. *Journ. Amer. Soc. Agron.* 22 (1930), 1004-1019.
- STRAUSS, J. L. and A. L. GRIZZARD : The effect of calcium, magnesium and potassium on peanut yields. *Soil Sci. Soc. of America, Proceedings* 1947, 12 (1948), 348-352.
- THORNTON, G. D. and F. E. BROADBENT : Preliminary greenhouse studies of the influence of nitrogen fertilization of peanuts on nodulation, yield and gynophore absorption of this element. *Journ. Amer. Soc. Agron.* 40 (1948), 64-69.
- WIERSUM, L. K. : Watertransport in the xylemas related to calcium uptake by groundnuts (*Arachis hypogaea* L.). *Plant and Soil* 3 (1951), 160-169.
- WOLK, P. C. VAN DER : Researches concerning geocarpy. *Publications sur la physiologie végétale*. Nimègue, 1914.

Erratum to

## INFLUENCE OF TILTH ON SOIL AND AIR-TEMPERATURE

R. H. A. VAN DUIN

Laboratory for Physics and Meteorology, Agricultural University,  
The Netherlands

In the article mentioned in the title (Neth. J. Agr. Sci. 2, 1954, 4 : 229-241) the phase shift between the heatflux into a layered soil and the temperature wave at the surface must read :

$$-\varphi = \arctg \frac{\frac{Q_s}{Q_a} \sin \left( \frac{\pi}{4} - \Delta \varphi_0 \right) + \sin (\arctg 1/2 j)}{\frac{Q_s}{Q_a} \cos \left( \frac{\pi}{4} - \Delta \varphi_0 \right) + \cos (\arctg 1/2 j)} \quad (17a)$$

where  $Q_s / Q_a$  is given by eq. 13a.

The results of the calculations in the case of a dry soil with  $q_s + q_a =$  constant are decreased by about 3% if eq. 17a is used. Owing to this correction the corresponding part of table 3 must be replaced by :

	z = 0				z = + 5 cm			
	I <sub>c</sub>	I <sub>d</sub>	II <sub>c</sub>	II <sub>d</sub>	I <sub>c</sub>	I <sub>d</sub>	II <sub>c</sub>	II <sub>d</sub>
A $T_{\min}$	18.3 - 4.3	21.6 - 7.6	27.8 (-18.8)	32.8 (-18.8)	14.6 - 0.6	17.3 (-3.3)	22.2 (-8.2)	26.2 (-12.2)